



## A genome-wide scan for common variants affecting the rate of age-related cognitive decline

Philip L. De Jager<sup>a,b,c,1,\*</sup>, Joshua M. Shulman<sup>a,b,c,1</sup>, Lori B. Chibnik<sup>a,b,c,1</sup>,  
 Brendan T. Keenan<sup>a,c</sup>, Towfique Raj<sup>a,b</sup>, Robert S. Wilson<sup>d</sup>, Lei Yu<sup>d</sup>, Sue E. Leurgans<sup>d</sup>,  
 Dong Tran<sup>a,b,c</sup>, Cristin Aubin<sup>a,b,c</sup>, Christopher D. Anderson<sup>c,e,f</sup>, Alessandro Biffi<sup>c,e,f</sup>,  
 Jason J. Corneveaux<sup>g</sup>, Matthew J. Huentelman<sup>g</sup>, Alzheimer's Disease Neuroimaging Initiative,  
 Jonathan Rosand<sup>b,c,e,f</sup>, Mark J. Daly<sup>b,c,e,f</sup>, Amanda J. Myers<sup>h,i</sup>, Eric M. Reiman<sup>g,k,1</sup>,  
 David A. Bennett<sup>d</sup>, Denis A. Evans<sup>d,1,\*</sup>

<sup>a</sup> Program in Translational NeuroPsychiatric Genomics, Institute for the Neurosciences, Departments of Neurology and Psychiatry, Brigham and Women's Hospital, Boston, MA, USA

<sup>b</sup> Harvard Medical School, Boston, MA, USA

<sup>c</sup> Program in Medical and Population Genetics, Broad Institute, Cambridge, MA, USA

<sup>d</sup> Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, USA

<sup>e</sup> Center for Human Genetic Research, Massachusetts General Hospital, Boston, MA, USA

<sup>f</sup> Department of Neurology, Massachusetts General Hospital, Boston, MA, USA

<sup>g</sup> Neurogenomics Division, Translational Genomics Research Institute and Arizona Alzheimer's Consortium, Phoenix, AZ, USA

<sup>h</sup> Department of Psychiatry and Behavioral Sciences, University of Miami, Miller School of Medicine, Miami, FL, USA

<sup>i</sup> Johnnie B. Byrd, Sr. Alzheimer's Center and Research Institute, Tampa, FL, USA

<sup>j</sup> Banner Alzheimer's Institute and Department of Psychiatry, University of Arizona, Phoenix, AZ, USA

<sup>k</sup> Department of Psychiatry, University of Arizona, Tucson, AZ, USA

<sup>l</sup> Rush University Medical Center, Rush Institute for Healthy Aging, Chicago, IL, USA

Received 2 June 2011; received in revised form 9 August 2011; accepted 16 September 2011

### Abstract

Age-related cognitive decline is likely promoted by accumulated brain injury due to chronic conditions of aging, including neurodegenerative and vascular disease. Because common neuronal mechanisms may mediate the adaptation to diverse cerebral insults, we hypothesized that susceptibility for age-related cognitive decline may be due in part to a shared genetic network. We have therefore performed a genome-wide association study using a quantitative measure of global cognitive decline slope, based on repeated measures of 17 cognitive tests in 749 subjects from the Religious Orders Study. Top results were evaluated in 3 independent replication cohorts, consisting of 2279 additional subjects with repeated cognitive testing. As expected, we find that the Alzheimer's disease (AD) susceptibility locus, *APOE*, is strongly associated with rate of cognitive decline ( $P_{DISC} = 5.6 \times 10^{-9}$ ;  $P_{JOINT} = 3.7 \times 10^{-27}$ ). We additionally discover a variant, *rs10808746*, which shows consistent effects in the replication cohorts and modestly improved evidence of association in the joint analysis ( $P_{DISC} = 6.7 \times 10^{-5}$ ;  $P_{REP} = 9.4 \times 10^{-3}$ ;  $P_{JOINT} = 2.3 \times 10^{-5}$ ). This variant influences the expression of 2 adjacent genes, *PDE7A* and *MTFR1*, which are potential regulators of inflammation and oxidative injury, respectively. Using aggregate measures of genetic risk, we find that known susceptibility loci for cardiovascular disease, type 2 diabetes, and inflammatory diseases are not significantly associated with cognitive decline in our cohort. Our results suggest that intermediate phenotypes, when coupled with larger sample sizes,

\* Corresponding author at: Program in Translational NeuroPsychiatric Genomics, Department of Neurology, Brigham and Women's Hospital, 77 Avenue Louis Pasteur, NRB 168C, Boston, MA 02115, USA. Tel.: +1 617 525 4529; fax: +1 617 525 5333.

E-mail address: pdejager@rics.bwh.harvard.edu (P.L. De Jager).

\* Alternate corresponding author at: Rush Institute for Healthy Aging, Rush University Medical Center, 1645 W Jackson Street, Suite 675, Chicago, IL 60612, USA. Tel.: +1 312 942 3350; fax: +1 312 942 2861.

E-mail address: Denis\_Evans@rush.edu (D.A. Evans).

<sup>1</sup> Contributed equally to this study.

may be a useful tool to dissect susceptibility loci for age-related cognitive decline and uncover shared molecular pathways with a role in neuronal injury.

© 2011 Elsevier Inc. All rights reserved.

**Keywords:** Cognitive decline; Genome-wide association study; Alzheimer's disease; Genetic risk score

## 1. Introduction

Decline in cognitive performance occurs with advancing age and is associated with a variety of common, age-related chronic medical conditions. Alzheimer's disease (AD) is the most prevalent cause of dementia (Reitz et al., 2011a); however, many other common adult illnesses, including type 2 diabetes (Croxson and Jagger, 1995; Grodstein et al., 2001; Reijmer et al., 2010), cerebrovascular disease (Desmond et al., 2000; Pendlebury and Rothwell, 2009), as well as other cardiovascular risk factors (Desmond et al., 1993; Warsch and Wright, 2010), and inflammatory disorders (Lucin and Wyss-Coray, 2009) have been implicated in age-related cognitive decline. Based on autopsy series from community-based cohorts, most individuals with dementia have multiple contributory pathologies at the time of death (Neuropathology Group of the Medical Research Council Cognitive Function and Ageing Study, 2001; Sonnen et al., 2007; Troncoso et al., 2008). It is likely that diverse forms of brain injury interact to accelerate cognitive decline. For example, it has been suggested that vascular-related brain injury may promote the development of AD pathology or, less directly, the clinical manifestation of AD-related cognitive decline (Launer et al., 2008; Schneider and Bennett, 2010; Warsch and Wright, 2010). Results from a variety of experimental paradigms for the study of neuronal injury and repair indicate that overlapping cellular and molecular mechanisms likely mediate the response to a diversity of central nervous system insults (Bishop et al., 2010; Cho et al., 2010; Lucin and Wyss-Coray, 2009; Martinez-Vicente and Cuervo, 2007; Ross and Poirier, 2004). Besides the local reactions to brain lesions, the ability of neuronal networks to adapt to and compensate for an accumulated burden of injury, sometimes referred to as cognitive reserve (Stern, 2009), likely has a substantial impact on the trajectory of cognitive decline. Studies of elder twins suggest substantial heritability in cognitive performance in late life (McClearn et al., 1997; Swan et al., 1990), and we hypothesize that a core genetic network might therefore impact susceptibility for rate of age-related cognitive decline.

Genome-wide association studies have proven a successful strategy for discovering susceptibility genes for complex human traits, including neurologic disorders, such as AD (Bertram and Tanzi, 2009). Besides the *apolipoprotein E* locus (*APOE*), these studies have identified common variants in *ABCA7*, *BIN1*, *CD2AP*, *CD33*, *CLU*, *CRI*, *EPHA1*, *MS4A4/MS4A6E*, and *PICALM* as associated with AD susceptibility (Harold et al., 2009; Hollingworth et al., 2011;

Lambert et al., 2009; Naj et al., 2011; Seshadri et al., 2010). While elucidating the functional impact of disease-associated genetic variants remains an active area of investigation, there is evidence that these genes may have important roles beyond AD pathogenesis in affecting other disorders potentially relevant to cognitive decline. For example, in addition to the well-known effect of the *APOE* locus in promoting AD risk, this locus has also been associated with dyslipidemia, cardiovascular disease, and increased cerebral infarcts (Eichner et al., 2002; Kim et al., 2003; McCarron et al., 1999). Similarly, polymorphisms in the *CRI* gene, encoding a complement receptor, have previously been associated with susceptibility for infectious disease, particularly malaria (Cockburn et al., 2004; Rowe et al., 1997). We have shown that polymorphisms in both *APOE* (Wilson et al., 2002a, 2002b) and *CRI* (Chibnik et al., 2011) have a measurable impact on age-related cognitive decline, including in subjects without dementia, and further, that these associations are mediated in part by an effect on promoting amyloid plaque pathology (Bennett et al., 2005a; Chibnik et al., 2011).

The Religious Orders Study (ROS) is following more than 1100 older Catholic nuns, priests, and brothers who have completed up to 16 years of annual cognitive testing. Here, we have leveraged available genotyping data for 749 subjects of European ancestry with longitudinal cognitive data to conduct a genome scan for loci associated with the rate of age-related cognitive decline. We report efforts to replicate the best results using data from 2 complementary, community-based studies, the Rush Memory and Aging Project (MAP) and Chicago Health and Aging Project (CHAP), as well as a predominantly clinic-derived subject sample from the Alzheimer's Disease Neuroimaging Initiative (ADNI), and offer evidence in support of replication for 1 variant. Finally, we explore whether known genetic susceptibility factors associated with other illnesses that are known to influence the risk of dementia, such as AD, cardiovascular disease, and type 2 diabetes, also affect age-related cognitive decline.

## 2. Methods

### 2.1. Subjects

Subjects are participants from 4 longitudinal studies, which are each described below. The number of study subjects with genotyping data, included in the genetic anal-

Table 1  
Demographic and clinical characteristics of study cohorts

	ROS	MAP	CHAP <sup>a</sup>	ADNI
<i>n</i>	749	825	737	717
Age at enrollment	75.3 (± 7.2)	80.8 (± 6.6)	72.0 (± 5.5)	75.3 (± 6.9)
Education (y)	18.2 (± 3.4)	14.8 (± 2.9)	14.6 (± 3.2)	15.6 (± 3.0)
Male	254 (34.0%)	222 (26.9%)	291 (39.5%)	422 (58.9%)
Cognitive decline slope	−0.007 (± 0.1)	0.012 (± 0.1)	0.001 (± 0.04)	−0.001 (± 0.04)
Cognitively normal <sup>b</sup>	444 (59.4%)	460 (55.8%)	265 (68.7%)	211 (29.4%)
Mild cognitive impairment <sup>b</sup>	151 (20.2%)	200 (24.2%)	71 (18.4%)	205 (28.6%)
Dementia <sup>b,c</sup>	152 (20.4%)	165 (20.0%)	46 (11.9%)	301 (42.0%)

Key: AD, Alzheimer's disease; ADNI, Alzheimer's Disease Neuroimaging Initiative; CHAP, Chicago Health and Aging Project; MAP, Memory and Aging Project; NINCDS, National Institute of Neurologic and Communicative Disorders and Stroke; ROS, Religious Orders Study.

<sup>a</sup> Diagnoses of dementia and mild cognitive impairment only available for a subset (*n* = 386) of CHAP.

<sup>b</sup> Cognitive status at time of last evaluation.

<sup>c</sup> Of the study cohorts, 19.7% of ROS, 19% of MAP, and 11% of CHAP met NINCDS criteria for possible or probable AD.

yses, are described in the Genotyping Methods subsection, and also summarized in Table 1.

The ROS, started in 1994, enrolls Catholic priests, nuns, and brothers, aged 53 or older from about 40 groups in 12 states. Since January 1994, 1132 participants completed their baseline evaluation, of whom 1001 are non-Hispanic white, and the follow-up rate of survivors exceeds 90%. Participants were free of known dementia at enrollment and agreed to annual clinical evaluations. More detailed description of the ROS can be found in prior publications (Bennett et al., 2006a).

The Rush Memory and Aging Project (MAP), started in 1997, enrolls older men and women from assisted living facilities in the Chicago area with no evidence of dementia at baseline. Since October 1997, 1285 participants completed their baseline evaluation, of whom 1118 were non-Hispanic white. The follow-up rate of survivors exceeds 90%. Similar to ROS, participants agreed to annual clinical evaluations. More detailed descriptions can be found in previous studies (Bennett et al., 2006a, 2005b).

The Chicago Health and Aging Project (CHAP) is a biracial (63% African American, 37% non-Hispanic European American) longitudinal population study of all participating residents 65 years of age and older of 4 adjacent neighborhoods of the south side of Chicago that examines risk factors for cognitive decline and AD. Participation was 78.6% of all community residents at baseline and 80% to 85% retention of survivors at follow-up. The study began in 1993 and enrolls successive age cohorts of community residents as they attain the age of 65 years. To date, 10,802 subjects have contributed data. The subjects included in these analyses are a stratified random sample of European American subjects. More details may be found in prior publications (Bienias et al., 2003; Evans et al., 2003).

The ADNI was launched in 2003 by the National Institute on Aging (NIA), the National Institute of Biomedical Imaging and Bioengineering (NIBIB), the Food and Drug Administration (FDA), private pharmaceutical companies, and nonprofit organizations, as a 5-year public-private part-

nership for the study of mild cognitive impairment (MCI) and early AD. The Principal Investigator of this initiative is Michael W. Weiner, MD, VA Medical Center and University of California, San Francisco. ADNI is the result of efforts of many coinvestigators from a broad range of academic institutions and private corporations, and subjects have been recruited from over 50 sites across the USA and Canada. The cohort includes 800 adults, ages 55 to 90, to participate in the research—approximately 200 cognitively normal older individuals to be followed for 3 years, 400 people with MCI to be followed for 3 years, and 200 people with early AD to be followed for 2 years. Data used in the preparation of this article were obtained from the ADNI database ([adni.loni.ucla.edu](http://adni.loni.ucla.edu)). For up-to-date information, see [www.adni-info.org](http://www.adni-info.org).

## 2.2. Clinical and cognitive evaluation

Supplementary Table 1 summarizes the specific longitudinal cognitive testing data that were available within our discovery and replication cohorts. The ROS and MAP studies annually administer 21 cognitive tests, of which 17 tests are in common. These 17 tests were incorporated into summary measures of 5 domains of cognitive function—episodic memory (7 tests), visuospatial ability (2 tests), perceptual speed (2 tests), semantic memory (3 tests), and working memory (3 tests)—as previously described (Bennett et al., 2002, 2005b; Wilson et al., 2002c, 2005). The tests from each area of cognition were converted to *z*-scores, using the mean and SD from the baseline evaluation of all participants, and averaged to yield summary measures of each area of cognitive function as previously described (Bennett et al., 2002, 2005b; Wilson et al., 2002c, 2005). The global cognition summary measure used in our primary analyses was computed by averaging the 5 summary scores for each cognitive subdomain. Summary measures have the advantage of minimizing floor and ceiling effects, and other sources of random variability. A valid summary score required that at least half of the component scores be present. The global cognition summary measures in the CHAP and

ADNI cohorts were constructed using identical procedures as ROS and MAP on the available longitudinal testing data (Supplementary Table 1). In CHAP, cognitive testing included a subset of 3 instruments from ROS and MAP. In ADNI, 8 tests overlapped with ROS and MAP, and we additionally incorporated available longitudinal data from 11 other tests.

The clinical diagnoses of dementia and AD were made following the recommendations of the joint working group of the National Institute of Neurologic and Communicative Disorders and Stroke and the AD and Related Disorders Association (McKhann et al., 1984), as previously described in detail (Bennett et al., 2006b). Mild cognitive impairment (MCI) referred to those individuals rated as cognitively impaired by the neuropsychologist but not demented by the examining physician, as previously described (Bennett et al., 2002).

### 2.3. Statistical modeling of cognitive decline

Mixed effects models were used to characterize individual paths of change in the cognitive summary measures, including terms for age, sex, and years of education as fixed effects (Laird and Ware, 1982; Wilson et al., 2000, 2002c). In this approach, each individual's path is assumed to follow the mean path of the group except for random effects that cause the initial level of function (i.e., intercept) to be higher or lower and the rate of change (i.e., slope) to be faster or slower. These random effects are assumed to follow a bivariate normal distribution. The random and fixed effects were then used to estimate individual trajectories of cognitive decline. Residual, individual cognitive decline slope terms were extracted from the mixed models, after adjustment for the effects of age, sex, and education. Person-specific, adjusted residual slopes were then used as a quantitative outcome phenotype for the genetic association analyses. These estimates equate to the difference between an individual's slope and the predicted slope of an individual of the same age, sex, and education level. This method, including adjustment for age, sex, and education was used for all 4 cohorts.

### 2.4. Genotyping

DNA from ROS and MAP subjects was extracted from whole blood, lymphocytes, or frozen postmortem brain tissue and genotyped on the Affymetrix GeneChip 6.0 (Santa Clara, CA, USA) platform at either the Broad Institute's Center for Genotyping ( $n = 1204$ ) or the Translational Genomics Research Institute ( $n = 674$ ). These 2 sets of data underwent the same quality control (QC) analysis in parallel, and genotypes were pooled. Only self-declared non-Hispanic Caucasians were genotyped to minimize population heterogeneity. The PLINK toolkit ([pnu.mgh.harvard.edu/~purcell/plink](http://pnu.mgh.harvard.edu/~purcell/plink); Purcell et al., 2007) was used to implement our QC pipeline. We applied standard quality control measures for subjects (genotype success rate  $> 95\%$ , gen-

otype-derived gender concordant with reported gender, excess inter/intraheterozygosity) and for single nucleotide polymorphisms (SNPs) (Hardy-Weinberg equilibrium  $p > 0.001$ ); minor allele frequency (MAF)  $> 0.01$ , genotype call rate  $> 0.95$ ; misshap test  $> 1 \times 10^{-9}$ ). Subsequently, EIGENSTRAT (Price et al., 2006) was used to identify and remove population outliers using default parameters. At the conclusion of the QC pipeline, data on 672,266 SNPs was available for 1709 total ROS and MAP subjects. A total of 749 ROS subjects and 825 MAP subjects with longitudinal cognitive data and high-quality genotyping data were available for the discovery and replication analyses, respectively (Table 1).

For the replication analysis, the top 50 independent SNPs ( $p < 10^{-4}$ , minor allele frequency  $> 10\%$ ) based on the ROS discovery stage analysis were extracted from the quality-controlled MAP genome-wide dataset (above). The generation and quality control procedures for the ADNI genotyping dataset was previously described (Biffi et al., 2010). A total of 717 ADNI subjects with longitudinal cognitive testing data and genotypes were available for our analyses. CHAP subject DNA was extracted from whole blood and genotyping of top SNPs from the discovery stage genome-wide scan using matrix-assisted laser desorption ionization time-of-flight mass spectrometry on a Sequenom MassARRAY platform (San Diego, CA, USA). After excluding subjects for failed genotyping exceeding the 10% threshold, 414 individuals remained for subsequent analysis (genotyping rate in these subjects was  $> 99\%$ ). All SNP allele frequencies satisfied Hardy-Weinberg equilibrium ( $p > 0.001$ ). For evaluation of AD susceptibility alleles and for the development of the aggregate genetic risk scores, SNP genotypes in ROS and MAP were imputed using MACH software (version 1.0.16a) (Scott et al., 2007) and HapMap release 22 CEU (build 36) (International HapMap Consortium et al., 2007) as a reference.

### 2.5. Genome-wide association analysis

The genome-wide association analysis in the ROS discovery cohort, as well as the targeted association analysis of the top 50 SNPs in the MAP, CHAP, and ADNI replication cohorts, were performed using linear regression implemented in PLINK software (Purcell et al., 2007). As described above, the outcome phenotype was the residual cognitive decline slope extracted from the mixed effects models, after adjustment for age, gender, and education. The association analysis in the ROS discovery cohort was additionally adjusted for the first 3 ancestry principal components calculated using EIGENSTRAT (Price et al., 2006). In order to perform a joint replication and study-wide meta-analysis, the cohort-specific PLINK association output was subsequently analyzed using METAL software (Willer et al., 2010). The default METAL parameters were used, in which meta-analysis is based on  $p$ -values and direction of effect, and



weighted by sample size in each cohort. The Manhattan plot was generated using Haploview software (Barrett et al., 2005), and the association plot was generated using the SNAP web-tool (Johnson et al., 2008).

### 2.6. Aggregate genetic risk scores

We developed 4 disease-specific cumulative genetic risk scores (GRSs) based on published genome wide significant ( $p < 10^{-8}$ ) SNPs for the following disease categories: Alzheimer's disease; cardiovascular disease (including myocardial infarction, low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, triglycerides, hypertension, and stroke); inflammatory disease (including celiac disease, Crohn's disease, irritable bowel disease, multiple sclerosis, psoriasis, rheumatoid arthritis, systemic lupus erythematosus, type 1 diabetes, and ulcerative colitis); and type 2 diabetes. Genome-wide significant SNPs for each disease category were identified through a literature review, primarily using the Catalogue of Published Genome-Wide Association Studies available online at [www.genome.gov/26525384](http://www.genome.gov/26525384). For the cardiovascular disease and inflammatory disease categories, which have multiple component outcomes, SNPs that had reference alleles which were both protective for a disease and a risk factor for another disease were removed. To limit the amount of correlation between SNPs, we further refined this list by identifying all pairs of SNPs with a linkage disequilibrium  $r^2 \geq 0.5$  and keeping only the SNP in those pairs with the lowest  $p$ -value. Once we identified a final SNP set, the 4 GRSs were created by summing up the number of category specific risk alleles for each individual. To assure that our scores were comprehensive, we used imputed allelic dosages for SNPs that were not genotyped within our population. We then examined the associations between the resulting genetic risk scores and global cognitive decline using linear regression models adjusted for age, sex, and education. The final list of SNPs and risk alleles included in each score, as well as relevant references, can be found in Supplementary Table 7.

### 2.7. Gene expression quantitative trait locus analysis

Gene expression levels were quantified using messenger RNA (mRNA) derived from peripheral blood mononuclear cells (PBMCs) of 228 subjects of European ancestry with relapsing-remitting (RR) multiple sclerosis (MS) using an Affymetrix Human Genome U133 plus 2.0. These data were collected between July 2002 and October 2007, as part of the Comprehensive Longitudinal Investigation of MS at the Brigham and Women's Hospital. The data are available on the Gene Expression Omnibus web site (GSE16214) (De Jager et al., 2009). DNA from each individual was genotyped on the Affymetrix GeneChip 6.0 platform as a part of a case-control multiple sclerosis meta-analysis (De Jager et al., 2009). Using a Spearman rank correlation, we tested for

association using an additive model for allelic dosage as an independent variable and residuals of expression as the dependent variable. Significance was established by comparing the association  $p$ -values to the empiric distribution of  $p$ -values generated by permuting expression phenotypes 10,000 times independently for each gene. Similar methods were used to evaluate expression within the publically available lymphoblastic cell line expression dataset from 60 CEU individuals in the HapMap project (Stranger et al., 2007).

## 3. Results

### 3.1. Characteristics of the discovery cohort

Following quality control, genome-wide genotype data (672,266 SNPs) were available on 749 non-Hispanic, white subjects from the ROS with longitudinal cognitive testing. Detailed cohort characteristics are presented in Table 1. The mean age at enrollment was 75 years, and subjects were followed for 9 years, on average (range 1–15 years of follow-up). Cognitive decline trajectories were quantified based on annual performance of 17 distinct neuropsychological tests sampling 5 cognitive domains (episodic memory, perceptual speed, semantic memory, visuospatial ability, and working memory) (Supplementary Table 1). As previously described (Wilson et al., 2002c), a subject's performance on each test was standardized and an average, aggregate measure of global cognitive performance was computed. At recruitment, all subjects were without known dementia. At their last evaluation, 59% of subjects retained normal cognition, 20% had mild cognitive impairment and 20% had a diagnosis of dementia. We used linear mixed effects modeling, including all available longitudinal cognitive testing, to obtain a residual cognitive decline slope for each individual, adjusting for age at enrollment, gender, and years of education (Fig. 1). This slope parameter describes the person-specific rate of global cognitive change as a quantitative, continuous outcome and is the primary outcome measure that we used in our association study.

### 3.2. Genome-wide association scan for age-related cognitive decline

We implemented our genome-wide association scan using the residual cognitive decline slope for each individual as an outcome trait (Fig. 2). The genomic inflation factor was 1.009, indicating no significant inflation of our test statistics. A selection of top results from the scan, based on our replication analysis, are shown in Table 2, and complete results ( $p < 10^{-4}$ ) are provided in Supplementary Tables 8 and 9. As expected from prior studies (Feskens et al., 1994; Haan et al., 1999; Henderson et al., 1995; Hyman et al., 1996; Jonker et al., 1998; McQueen et al., 2007; Wilson et al., 2002a, 2002b), the strongest associations with the rate of cognitive decline were found for markers at the *APOE*




Fig. 1. Variability in the rate of age-related cognitive decline. Linear cognitive trajectories are shown for 100 random subjects from the Religious Orders Study (ROS) cohort, based on mixed effect modeling of repeated measures of the global cognition summary score, incorporating 17 distinct cognitive tests. Trajectories are adjusted for the effects of baseline age, gender, and education. The residual cognitive decline slope was used as an outcome for the genome-wide association analysis. The distribution of the cognitive decline trait for the entire ROS discovery cohort is shown in Supplementary Fig 1.

locus. No other locus association surpassed the genome-wide significance threshold ( $p < 5 \times 10^{-8}$ ). However, numerous polymorphisms demonstrate suggestive evidence of association ( $p < 10^{-4}$ ) with rate of cognitive decline, including several that fall within or adjacent to candidate genes previously implicated in cognition (*CTNND2*, *rs2973488*,  $p = 1.8 \times 10^{-7}$ ) (Israely et al., 2004; Medina et al., 2000) or AD susceptibility (*SORCS1*, *rs12219216*,  $p = 8.0 \times 10^{-5}$ ) (Reitz et al., 2011b). As we have previously reported (Chibnik et al., 2011), we find nominal evidence of association at the *CRI* locus (*rs6656401*,  $p = 0.048$ ), but do not detect evidence of association for other known AD susceptibility variants with the rate of cognitive decline in our discovery cohort (Supplementary Table 2).

### 3.3. Replication of variants associated with rate of cognitive decline

To replicate the results of our genome scan, the top 50 independent SNPs ( $p < 10^{-4}$ , minor allele frequency  $> 10\%$ ) from our discovery stage were evaluated in 3 additional cohorts of older individuals with longitudinal measures of cognitive performance (Supplementary Tables 2 and 8). MAP, CHAP, and ADNI are described in the Methods and the cohort characteristics are summarized in Table 1. A summary of all cognitive testing data used from each study is provided in Supplementary Table 1. As in ROS, available cognitive data for MAP, CHAP, and ADNI subjects were incorporated into aggregate global cognition scores, and mixed effect modeling of longitudinal data were used to compute residual, adjusted global cognitive decline slopes for all subjects. The SNPs selected for replication were evaluated in each of the 3 replication cohorts, and meta-analysis was implemented to compute summary association statistics ( $P_{REP}$ ) across the replication cohorts, with a total of 2279 subjects. We also determined a study-wide association meta-analysis statistic ( $P_{JOINT}$ ), consisting of 3028 total subjects, including the results from the discovery cohort and those of the 3 replication cohorts. The results of the replication and joint analyses are shown in Table 2. We hypothesized that discovered variants that truly impact rate of cognitive decline would show robust and consistent effects across these sample collections, despite modest differences in cohort make-up, such as mean age, cognitive testing procedures, and years of follow-up.

As expected, a SNP at the *APOE* locus, *rs4420638*, showed significant replication for association with rate of cognitive decline across each independent sample evaluated and in the pooled replication cohort ( $P_{MAP} = 9.4 \times 10^{-9}$ ,  $P_{CHAP} = 5.1 \times 10^{-6}$ ,  $P_{ADNI} = 5.3 \times 10^{-8}$ ,  $P_{REP} = 9.1 \times 10^{-20}$ ), and the overall association was strongly enhanced in the joint analysis ( $P_{ROS} = 5.6 \times 10^{-9}$ ,  $P_{JOINT} = 3.7 \times$




Fig. 2. A genome-wide association scan for age-related cognitive decline. Using the residual cognitive decline slope as an outcome trait, associations were evaluated for 672,266 single nucleotide polymorphisms (SNPs) in the discovery cohort consisting of 749 Religious Orders Study (ROS) subjects. (A) Quantile-quantile plot. (B) Manhattan plot. Thresholds for suggestive ( $p < 10^{-4}$ , blue) and genome-wide ( $p < 5 \times 10^{-8}$ , red) significance are indicated.

Table 2

Top results of genome-wide scan for rate of cognitive decline

SNP	C	A1/2	MAF	Discovery (ROS)		Replication				Joint	Direction	Gene(s)
				$\beta$ (95% CI)	$p$	$p_{MAP}$	$p_{CHAP}$	$p_{ADNI}$	$p_{REP}$	$p$		
rs4420638	19	C/T	0.19	-0.039 (-0.051 to -0.026)	$5.62 \times 10^{-9}$	$9.38 \times 10^{-9}$	$5.06 \times 10^{-6}$	$5.31 \times 10^{-8}$	$9.12 \times 10^{-20}$	$3.74 \times 10^{-27}$	++++	TOMM40, APOE
rs10808746	8	T/C	0.43	0.021 (0.011 to 0.031)	$6.73 \times 10^{-5}$	0.091	0.274	0.089	$9.44 \times 10^{-3}$	$2.30 \times 10^{-5}$	++++	PDE7A, MTFR1
rs10769565	11	G/A	0.34	-0.023 (-0.034 to -0.013)	$9.09 \times 10^{-6}$	0.455	0.159	NA	0.131	$1.63 \times 10^{-4}$	++++?	OR56A4, OR56A1
rs9387454	6	A/G	0.36	-0.023 (-0.033 to -0.012)	$2.12 \times 10^{-5}$	0.870	0.014	0.710	0.199	$1.26 \times 10^{-3}$	----+	
rs11939527	4	T/C	0.25	-0.023 (-0.034 to -0.011)	$9.55 \times 10^{-5}$	0.876	0.196	0.154	0.149	$1.41 \times 10^{-3}$	+-+-	EVC
rs376535	2	C/T	0.10	-0.034 (-0.050 to -0.018)	$4.38 \times 10^{-5}$	0.113	0.031	0.097	0.230	$2.10 \times 10^{-3}$	+-+-	SLC8A1
rs2571577	6	C/T	0.20	-0.027 (-0.039 to -0.014)	$3.42 \times 10^{-5}$	0.688	0.039	0.715	0.255	$2.32 \times 10^{-3}$	+-+-	LAMA2
rs13015892	2	A/G	0.17	-0.031 (-0.044 to -0.017)	$1.50 \times 10^{-5}$	0.790	0.589	0.162	0.347	$3.04 \times 10^{-3}$	+-+-	
rs9602785	13	C/T	0.15	-0.033 (-0.047 to -0.019)	$5.71 \times 10^{-6}$	0.478	0.770	0.843	0.482	$4.17 \times 10^{-3}$	++++	SLITRK6
rs11701130	21	A/C	0.26	0.024 (0.013 to 0.036)	$5.06 \times 10^{-5}$	0.141	0.101	0.116	0.349	$4.67 \times 10^{-3}$	+-+-	PCBP3
rs4760608	12	C/A	0.21	-0.027 (-0.040 to -0.015)	$2.68 \times 10^{-5}$	0.674	0.100	0.779	0.398	$4.77 \times 10^{-3}$	+-+-	COL2A1
rs2973488	5	T/A	0.11	-0.044 (-0.061 to -0.028)	$1.81 \times 10^{-7}$	0.823	0.688	0.524	0.792	$5.06 \times 10^{-3}$	+-+-	CTNND2
rs7606972	2	G/T	0.36	0.022 (0.012 to 0.033)	$2.08 \times 10^{-5}$	0.541	0.803	0.811	0.520	$7.57 \times 10^{-3}$	—	
rs9875783	3	A/C	0.25	-0.024 (-0.036 to -0.012)	$5.78 \times 10^{-5}$	0.894	0.803	0.075	0.437	$7.62 \times 10^{-3}$	+-+-	
rs524398	2	A/G	0.42	-0.022 (-0.032 to -0.012)	$2.45 \times 10^{-5}$	0.155	0.971	0.736	0.519	$7.84 \times 10^{-3}$	+-+-	THSD7B
rs610315	10	A/T	0.26	-0.024 (-0.036 to -0.013)	$3.00 \times 10^{-5}$	0.634	0.090	0.918	0.535	$8.94 \times 10^{-3}$	+-+-	AFAP1L2
rs6919160	6	A/T	0.31	-0.024 (-0.035 to -0.013)	$1.72 \times 10^{-5}$	0.562	0.354	0.521	0.589	$9.14 \times 10^{-3}$	+-+-	
rs347311	1	C/G	0.31	-0.025 (-0.036 to -0.014)	$7.62 \times 10^{-6}$	0.446	0.094	0.791	0.729	0.012	+-+-	NOS1AP
rs4129955	4	T/C	0.47	-0.020 (-0.030 to -0.010)	$6.69 \times 10^{-5}$	0.195	0.589	0.123	0.693	0.020	+-+-	
rs3212701	19	A/G	0.25	0.052 (0.028 to 0.075)	$2.69 \times 10^{-5}$	0.761	0.661	0.397	0.456	0.021	+-+-	JAK3
rs1595781	3	G/A	0.31	-0.023 (-0.034 to -0.012)	$6.70 \times 10^{-5}$	0.665	0.617	0.551	0.717	0.022	+-+-	
rs276420	13	A/C	0.45	-0.021 (-0.031 to -0.011)	$3.25 \times 10^{-5}$	0.226	0.947	0.266	0.887	0.028	----	
rs12660119	6	G/A	0.11	-0.038 (-0.054 to -0.021)	$7.28 \times 10^{-6}$	0.912	0.792	0.827	0.970	0.029	+-+-	WASF1
rs1041676	1	T/C	0.32	-0.023 (-0.034 to -0.012)	$4.73 \times 10^{-5}$	0.567	0.998	0.384	0.882	0.032	+-+-	NEGR1
rs2086366	15	T/C	0.37	0.020 (0.010 to 0.030)	$4.77 \times 10^{-5}$	0.992	0.392	0.827	0.706	0.039	+-+-	
rs12219216	10	A/G	0.41	0.020 (0.010 to 0.031)	$8.01 \times 10^{-5}$	0.777	0.359	0.415	0.915	0.040	+-+-	SORCS1
rs1534378	7	T/C	0.11	-0.030 (-0.044 to -0.015)	$5.51 \times 10^{-5}$	0.848	0.729	0.654	0.714	0.042	+-+-	
rs12362733	11	C/T	0.19	0.027 (0.015 to 0.040)	$2.72 \times 10^{-5}$	0.267	0.168	0.643	0.891	0.049	+-+-	LUZP2

Summary results are shown for single nucleotide polymorphisms (SNPs) with the best evidence of association in the study-wide meta-analysis ( $p_{JOINT} < 0.05$ ). The top 50 independent SNPs from the Religious Orders Study (ROS) discovery cohort (minor allele frequency [MAF] > 10% and LD-pruned based on  $r^2 > 0.5$ ) were evaluated for replication (complete results presented in Supplementary Table 8).  $\beta$  (95% CI) and  $p$ -values are shown for association with rate of cognitive decline in the discovery stage.  $p$ -values are also shown for associations in the memory and aging project (MAP), Chicago Health and Aging Project (CHAP), and ADNI replication cohorts. Meta-analytic  $p$ -values are shown for the replication cohorts ( $p_{REP}$ ) and for the study-wide analysis including all cohorts ( $p_{JOINT}$ ). Direction of effect (+ = protective, - = risk) is indicated for all 4 cohorts.

Key: A1/2, minor allele/major allele; ADNI, Alzheimer's Disease Neuroimaging Initiative; C, chromosome; CI, confidence interval; LD, linkage disequilibrium; MAF, minor allele frequency in ROS cohort; NA, not applicable; ROS, Religious Orders Study.




Fig. 3. Association of *rs10808746* at the *PDE7A/MTFR1* locus with rate of cognitive decline. (A) Mean linear trajectories of cognitive decline within the Religious Orders Study (ROS) discovery cohort for each of the *rs10808746* genotype classes, demonstrating evidence that the *rs10808746<sup>G</sup>* allele is associated with increased rate of cognitive decline. (B) Plot showing *rs10808746* association peak over the *PDE7A*, *MTFR1*, and *ARMCI* genes.

$10^{-27}$ ). Although no other variant met a threshold of genome-wide significance at the conclusion of the replication study, a chromosome 8 SNP, *rs10808746*, showed consistent direction of effect on the rate of cognitive decline in each cohort (the major allele *rs10808746<sup>G</sup>* is associated with increased risk of cognitive decline) and suggestive evidence of replication in the combined replication cohort ( $P_{REP} = 0.009$ ). Further supporting replication, in the study-wide joint analysis, *rs10808746* shows modest evidence of enhanced association when all available data are considered ( $P_{ROS} = 6.7 \times 10^{-5}$ ,  $P_{JOINT} = 2.3 \times 10^{-5}$ ).

Figure 3A shows the mean trajectory of cognitive decline within the discovery cohort for the 3 genotype classes of *rs10808746*. These data are supportive of the major allele *rs10808746<sup>G</sup>* being associated with more rapid cognitive decline. To better understand the impact of this variant on cognition, we evaluated associations with cognitive decline based on measures for each of the 5 domains of the ROS global cognitive score (Supplementary Table 3). This polymorphism was associated with the rate of cognitive decline in episodic memory ( $p = 3 \times 10^{-4}$ ), perceptual speed ( $p = 0.048$ ), semantic memory ( $p = 0.015$ ), and working memory ( $p = 0.0046$ ), but not visuospatial processing ( $p = 0.13$ ), suggesting that, while it appears to have a predominant effect on decline in episodic memory, its role may not be limited to a single, functionally distinct anatomic region or circuit.

#### 3.4. Analysis of *rs10808746* effect on local gene expression

The *rs10808746* SNP falls within an intron of phosphodiesterase 7A (*PDE7A*) and is near 2 flanking genes, mitochondrial fission regulator 1 (*MTFR1*) and armadillo repeat-containing protein 1 (*ARMCI*), that fall within a linkage disequilibrium block identified by the association peak (Fig. 3B). In order to begin to characterize the effect of the discovered variant, we attempted to evaluate its impact

on local gene expression in available datasets. In an analysis of gene expression data from PBMCs of 228 individuals with demyelinating disease (De Jager et al., 2009), representing a set of subjects with an activated immune system, we found evidence for association between the discovered variant and expression of both *PDE7A* ( $p = 8.4 \times 10^{-4}$ ) and *MTFR1* ( $p = 4.5 \times 10^{-5}$ ), but not *ARMCI* ( $p = 0.21$ ) (Fig. 4, Supplementary Table 4). The significance of the *rs10808746* association with *PDE7A* and *MTFR1* expression in PBMCs was robust to gene-based permutation testing. Published data from a different cell type, HapMap lymphoblastic cell lines (Stranger et al., 2007), also revealed associations with the expression of *PDE7A* ( $p = 5 \times 10^{-3}$ ) and *MTFR1* ( $p = 0.041$ ), but not *ARMCI* ( $p = 0.17$ ). However, the direction of effect for the association between this SNP and gene expression was not consistent between these 2 sets of data generated from different cell populations: the risk allele was associated with decreased expression of both genes in the PBMCs from subjects with demyelinating disease, but a more modest increase in both *PDE7A* and *MTFR1* expression in the HapMap cell lines, which are B cells transformed by Epstein-Barr virus. Based on available databases of gene expression data, such as BioGPS (biogps.gnf.org), both *PDE7A* and *MTFR1* are widely expressed in diverse tissue types, including the central nervous system (Rhead et al., 2010; Su et al., 2002; Wu et al., 2009); however, a publicly-available brain expression dataset (Myers et al., 2007) with genome-wide genotyping did not include probes for either *MTFR1* or *PDE7A*. We were therefore unable to readily confirm our findings in this tissue context.

#### 3.5. Aggregate genetic risk profiles for prediction of cognitive decline

In addition to a per SNP genome-wide analysis, we pursued a complementary approach taking into account the aggregate effect of multiple validated susceptibility alleles



Fig. 4. Association of *rs10808746* with *MTFR1* and *PDE7A* gene expression. The relation of *rs10808746* genotype was evaluated with locus transcript levels in an available gene expression dataset from 228 subjects with demyelinating disease. The *rs10808746<sup>G</sup>* allele is associated with decreased expression of *MTFR1* ( $p = 4.5 \times 10^{-5}$ ) and *PDE7A* ( $p = 8.4 \times 10^{-4}$ ), but not *ARMCI* ( $p = 0.21$ ).

for other diseases. Specifically, we developed 4 separate genetic risk score models incorporating known susceptibility variants for medical conditions known or hypothesized to promote cognitive decline, including AD, cardiovascular disease, type 2 diabetes, and inflammatory disease. In our ROS discovery cohort, a diagnosis of AD or of type 2 diabetes mellitus was significantly associated with cognitive decline (Supplementary Table 5), consistent with prior studies (Arvanitakis et al., 2004). Supplementary Table 7 summarizes the SNPs included in each aggregate genetic risk score, along with the relevant references. For both the AD and cardiovascular risk models, we excluded *APOE* from consideration, because this locus' strong effect on cognitive decline would overwhelm the contribution of loci with more modest effect sizes. As shown in Table 3, none of the aggregate risk models returned evidence of association in our ROS discovery cohort, and only the AD model demonstrated nominal association to cognitive decline in a more powerful joint analysis that included both the ROS and MAP cohorts. Our modest sample size limits our power in these analyses, but validating our strategy, we do see the expected correlation of the aggregate estimate of genetic

risk for type 2 diabetes with a diagnosis of diabetes in our subjects (Supplementary Table 6).

### 3.6. Power for discovery and replication of genetic variants associated with cognitive decline

The genetic architecture of cognitive decline is currently not known. Given our results and the size of our discovery and replication sample sets, it is likely that cognitive decline, like most other complex traits, is influenced by many variants of modest effect, in addition to the *APOE* locus which has a strong effect on this trait. Using the *PDE7A/MTFR1* and *CRI* loci as putative susceptibility loci with which to calibrate the design of future studies exploring cognitive decline, we performed power calculations to assess the sample size that would be necessary for a robust discovery study. For *CRI*, we observe an effect size ( $\beta$ ) of  $-0.0130$  and a MAF of 0.20 in the ROS cohort; thus, we would need 7241 subjects to observe a genome-wide significant effect on the rate of cognitive decline ( $p < 5 \times 10^{-8}$ ) with 80% power, and 8288 subjects for 90% power. For *PDE7A/MTFR1*, the effect size in the ROS cohort ( $\beta =$

Table 3  
Associations between aggregate genetic risk scores and rate of cognitive decline

GRS	ROS		MAP		ROS + MAP	
	$\beta$ (95% CI)	$p$	$\beta$ (95% CI)	$p$	$\beta$ (95% CI)	$p$
AD	-0.001 (-0.005 to 0.003)	0.50	-0.004 (-0.008 to 0.000)	0.06	-0.003 (-0.005 to 0.000)	0.07
CVD	-0.0004 (-0.002 to 0.001)	0.61	0.001 (-0.001 to 0.002)	0.49	0.0001 (-0.001 to 0.001)	0.87
I	0.0003 (-0.0006 to 0.001)	0.49	0.0001 (-0.0008 to 0.001)	0.82	0.0002 (-0.0004 to 0.001)	0.52
T2D	-0.001 (-0.003 to 0.001)	0.28	-0.0001 (-0.002 to 0.002)	0.96	-0.001 (-0.002 to 0.001)	0.48

$\beta$  for association of aggregate genetic risk score (GRS) and rate of cognitive decline. Separate GRS were developed based on published single nucleotide polymorphism (SNP) associations with Alzheimer's disease (AD), cardiovascular disease (CVD), inflammatory disease (I), and type 2 diabetes (T2D).

Key: CI, confidence interval; MAP, Memory and Aging Project; ROS, Religious Orders Study.

0.0210) is probably inflated given that this is the discovery cohort for this variant. However, given this magnitude of an effect and a MAF of 0.43, we would need 1794 subjects to have 80% power to observe a genome-wide significant effect ( $p < 5 \times 10^{-8}$ ), and 2053 subjects for 90% power. However, in the MAP cohort, the *PDE7A/MTFR1* has an effect size of 0.008, yielding an alternative estimate of 12,175 subjects needed for 80% power and 13,937 subjects for 90% power. The latter estimate is likely to more accurately reflect the true effect size of our variant because it comes from a replication cohort. The requirement for a larger sample size is consistent with what has been observed for other human traits (Park et al., 2010) and likely explains, at least in part, why we have not yet discovered susceptibility variants for cognitive decline reaching the genome-wide significance level. These estimates will guide the design of our future efforts: it appears that sample sizes between 7241 and 12,175 subjects will be needed to have reasonable power to assemble convincing evidence of association between genetic variation and the rate of cognitive decline.

#### 4. Discussion

We report the results of a genome-wide scan in 749 elder subjects to identify loci associated with the rate of cognitive decline, using a mixed effects model that incorporates repeated cognitive measures. Consistent with numerous prior studies (Feskens et al., 1994; Haan et al., 1999; Henderson et al., 1995; Hyman et al., 1996; Jonker et al., 1998; Wilson et al., 2002a, 2002b), we found robust evidence for association between the *APOE* locus and the rate of age-related cognitive decline. Besides *APOE*, in a replication analysis including 2279 additional subjects, we identified 1 other locus (*PDE7A/MTFR1*) with consistent direction of effect on cognitive decline in all cohorts and improved evidence of association in a joint analysis. While additional replication efforts will be needed to validate the role of this locus in cognitive decline, our replication and extension analyses strongly suggest that this locus has an effect on the rate of cognitive decline in aging individuals.

To begin to understand the function of this variant, we explored its possible role in influencing the expression of nearby genes using available gene expression data sets. Although further work is needed to better understand the effect of this variant in different cell types (particularly in the brain for which data were not available), 2 of the 3 genes found in the block of LD surrounding the associated variant—*PDE7A* and *MTFR1*—appear to be influenced by *rs10808746*, which is found in an intron of *PDE7A*. *MTFR1* has been shown to induce mitochondrial fission in a variety of cell types (Tonachini et al., 2004), and in *Mtfr1*<sup>null</sup> mice, testicular cells show increased oxidative injury and reduced expression of free radical scavengers, such as glutathione peroxidase 3 (Monticone et al., 2007). Thus, *MTFR1* is a

reasonable candidate gene for further validation in the context of cognitive decline because it could be involved in modulating the response of central nervous system (CNS) cells to oxidative stress, implicated in many different models of neuronal injury including AD, amyotrophic lateral sclerosis, and Parkinson's disease (Cho et al., 2010). *PDE7A* is another strong candidate: it is expressed in both neuronal and nonneuronal CNS cells and regulates intracellular signaling by affecting the concentration of cyclic dinucleotides (Miró et al., 2001; Pérez-Torres et al., 2003). Data on its possible function in CNS cells is limited, but 1 report suggests that inhibiting this molecule may contribute to enhancing cell death in response to an apoptotic signal in lymphocytes (Dong et al., 2010). There is also mixed evidence that it may have a role in mediating the expression of proinflammatory cytokines during T cell activation (Kadoshima-Yamaoka et al., 2009; Yang et al., 2003). Thus, *PDE7A* is also a reasonable candidate for mediating the effect of the *rs10808746* variant, and it is further possible that the variant influences the expression and function of both genes, possibly by affecting shared pathways involved in neuronal degeneration.

Based on our power calculations, it is clear that substantially larger sample sizes including 10,000–15,000 subjects will be needed for a definitive investigation of the genetic architecture of the rate of cognitive decline. This estimate is in line with power calculations for other complex traits (Park et al., 2010). The distribution of results from the ROS cohort suggest that it is unlikely that loci with a strong effect on cognitive decline exist outside of *APOE*. The existence of loci with more modest effects on cognitive decline is consistent with the suggestive evidence presented in support of *CRI*, a known AD susceptibility locus, and possibly *PDE7A/MTFR1*. We attempted to enhance our ability to test detect the role of genetic variation in cognitive decline by assessing aggregate measures of genetic risk for conditions that are well-known (AD, cerebrovascular disease, and type 2 diabetes) or hypothesized (inflammatory diseases) to influence this trait. Our aggregate measure of AD risk is nominally associated with cognitive decline, suggesting that this may be a fruitful strategy to deploy in future studies that leverage larger sample sizes and additional susceptibility alleles that will emerge in the near future.

Our study was based on the hypothesis that common genetic pathways mediate the response and adaptation to diverse forms of brain pathology, and that variation in such loci might therefore impact age-related cognitive decline. This hypothesis is supported by observations from a variety of experimental paradigms of neuronal injury that several core cellular pathways, including protein misfolding/aggregation (Ross and Poirier, 2004), autophagy (Martinez-Vicente and Cuervo, 2007), mitochondrial dynamics (Cho et al., 2010), and inflammation (Lucin and Wyss-Coray, 2009), play important roles in multiple neurodegenerative diseases. In prospective, community-based autopsy series of

dementia, most brains are found to have multiple pathologies (Neuropathology Group of the Medical Research Council Cognitive Function and Ageing Study, 2001; Schneider et al., 2007; Sonnen et al., 2007), which likely interact to produce the clinical manifestations of age-related cognitive decline (Wilson et al., 2010). Our strategy of using cognitive trajectories as an outcome trait for a genome-wide scan holds great promise for the identification of core genetic mechanisms mediating cognitive reserve and adaptation to injury. However, our current study provides evidence that the genetic architecture of this trait may be similar to that of other complex human traits, involving common variants of modest effects. We provide estimates of the substantially larger sample size required to achieve effective statistical power for robust gene discovery efforts, highlighting the need to combine additional cohorts with those reported here.

### Disclosure statement

The authors have no conflicts of interest to report related to this work.

### Acknowledgements

The authors thank Sara Pulit for help with assembling a comprehensive list of SNPs associated with inflammatory diseases. The authors are also grateful to the participants in the Religious Orders Study, the Chicago Health and Aging Project, the Memory and Aging Project, and the Alzheimer's Disease Neuroimaging Initiative. This work is supported by the National Institutes of Health [R01 AG030146, P30 AG10161, R01 AG17917, R01 AG15819, K08 AG034290, P30 AG10161 and R01 AG11101], and the Illinois Department of Public Health. J.M.S. is additionally supported by the Clinical Investigator Training Program: Beth Israel Deaconess Medical Center and Harvard/MIT Health Sciences and Technology, in collaboration with Pfizer Inc. and Merck & Co. A.J.M. is supported by NIA grant R01 AG034504 and the Johnnie B Byrd Institute. M.J.H., J.C., and E.M.R. are supported by grants from Kronos Life Sciences Laboratories, the National Institute on Aging (Arizona Alzheimer's Disease Center P30 AG19610, R01 AG023193, Mayo Clinic Alzheimer's Disease Center P50 AG16574 and Intramural Research Program), the National Alzheimer's Coordinating Center (U01 AG016976), and the state of Arizona. Additional funding was obtained from the NIH Neuroscience Blueprint (U24NS051872), the ENDGAME Consortium (U01HL084744), a National Institute on Aging grant to Carl Cotman (University of California, Irvine, P50 AG23173), National Institute on Aging (K01AG024079) and the state of Arizona. Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health grant U01 AG024904). ADNI is funded by

the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Abbott, AstraZeneca AB, Bayer Schering Pharma AG, Bristol-Myers Squibb, Eisai Global Clinical Development, Elan Corporation, Genentech, GE Healthcare, GlaxoSmithKline, Innogenetics, Johnson and Johnson, Eli Lilly and Co., Medpace, Inc., Merck and Co., Inc., Novartis AG, Pfizer Inc, F. Hoffman-La Roche, Schering-Plough, Synarc, Inc., as well as nonprofit partners the Alzheimer's Association and Alzheimer's Drug Discovery Foundation, with participation from the US Food and Drug Administration. Private sector contributions to ADNI are facilitated by the Foundation for the National Institutes of Health ([www.fnih.org](http://www.fnih.org)). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of California, Los Angeles. This research was also supported by NIH grants; 1P30 AG010129, K01 AG030514, and the Dana Foundation.

Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database ([adni.loni.ucla.edu](http://adni.loni.ucla.edu)). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A full listing of ADNI investigators can be found at: [adni.loni.ucla.edu/wpcontent/uploads/how\\_to\\_apply/ADNI\\_Authorship\\_List.pdf](http://adni.loni.ucla.edu/wpcontent/uploads/how_to_apply/ADNI_Authorship_List.pdf).

### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.neurobiolaging.2011.09.033](https://doi.org/10.1016/j.neurobiolaging.2011.09.033).

### References

- Arvanitakis, Z., Wilson, R.S., Bienias, J.L., Evans, D.A., Bennett, D.A., 2004. Diabetes mellitus and risk of Alzheimer disease and decline in cognitive function. *Arch. Neurol.* 61, 661–666.
- Barrett, J.C., Fry, B., Maller, J., Daly, M.J., 2005. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21, 263–265.
- Bennett, D., Schneider, J., Arvanitakis, Z., Kelly, J., Aggarwal, N., Shah, R., Wilson, R., 2006a. Neuropathology of older persons without cognitive impairment from two community-based studies. *Neurology* 66, 1837–1844.
- Bennett, D., Schneider, J., Wilson, R., Bienias, J., Berry-Kravis, E., Arnold, S., 2005a. Amyloid mediates the association of apolipoprotein E e4 allele to cognitive function in older people. *J. Neurol. Neurosurg. Psychiatry* 76, 1194–1199.
- Bennett, D.A., Schneider, J.A., Aggarwal, N.T., Arvanitakis, Z., Shah, R.C., Kelly, J.F., Fox, J.H., Cochran, E.J., Arends, D., Treinkman, A.D., Wilson, R.S., 2006b. Decision rules guiding the clinical diagnosis of Alzheimer's disease in two community-based cohort studies compared to standard practice in a clinic-based cohort study. *Neuroepidemiology* 27, 169–176.

- Bennett, D.A., Schneider, J.A., Buchman, A.S., Mendes de Leon, C., Bienias, J.L., Wilson, R.S., 2005b. The Rush Memory and Aging Project: study design and baseline characteristics of the study cohort. *Neuroepidemiology* 25, 163–175.
- Bennett, D.A., Wilson, R.S., Schneider, J.A., Evans, D.A., Beckett, L.A., Aggarwal, N.T., Barnes, L.L., Fox, J.H., Bach, J., 2002. Natural history of mild cognitive impairment in older persons. *Neurology* 59, 198–205.
- Bertram, L., Tanzi, R.E., 2009. Genome-wide association studies in Alzheimer's disease. *Hum. Mol. Genet.* 18, R137–R145.
- Bienias, J.L., Beckett, L.A., Bennett, D.A., Wilson, R.S., Evans, D.A., 2003. Design of the Chicago Health and Aging Project (CHAP). *J. Alzheimers Dis.* 5, 349–355.
- Biffi, A., Anderson, C.D., Desikan, R.S., Sabuncu, M., Cortellini, L., Schmansky, N., Salat, D., Rosand, J., Alzheimer's Disease Neuroimaging Initiative (ADNI), 2010. Genetic variation and neuroimaging measures in Alzheimer disease. *Arch. Neurol.* 67, 677–685.
- Bishop, N.A., Lu, T., Yankner, B.A., 2010. Neural mechanisms of ageing and cognitive decline. *Nature* 464, 529–535.
- Chibnik, L.B., Shulman, J.M., Leurgans, S.E., Schneider, J.A., Wilson, R.S., Tran, D., Aubin, C., Buchman, A.S., Heward, C.B., Myers, A.J., Hardy, J.A., Huentelman, M.J., Corneveaux, J.J., Reiman, E.M., Evans, D.A., Bennett, D.A., De Jager, P.L., 2011. CR1 is associated with amyloid plaque burden and age-related cognitive decline. *Ann. Neurol.* 69, 560–569.
- Cho, D.H., Nakamura, T., Lipton, S.A., 2010. Mitochondrial dynamics in cell death and neurodegeneration. *Cell. Mol. Life Sci.* 67, 3435–3447.
- Cockburn, I.A., Mackinnon, M.J., O'Donnell, A., Allen, S.J., Moulds, J.M., Baisor, M., Bockarie, M., Reeder, J.C., Rowe, J.A., 2004. A human complement receptor 1 polymorphism that reduces Plasmodium falciparum rosetting confers protection against severe malaria. *Proc. Natl. Acad. Sci. U. S. A.* 101, 272–277.
- Crosson, S.C., Jagger, C., 1995. Diabetes and cognitive impairment: a community-based study of elderly subjects. *Age Ageing* 24, 421–424.
- De Jager, P., Jia, X., Wang, J., de Bakker, P., Ottoboni, L., Aggarwal, N., Piccio, L., Raychaudhuri, S., Tran, D., Aubin, C., Briskin, R., Romano, S., Consortium, I.M.G., Baranzini, S., McCauley, J., Pericak-Vance, M., Haines, J., Gibson, R., Naeglin, Y., Uitdehaag, B., Matthews, P., Kappos, L., Polman, C., McArdle, W., Strachan, D., Evans, D., Cross, A., Daly, M., Compston, A., Sawcer, S., Weiner, H., Hauser, S., Hafler, D., Oksenberg, J., 2009. Meta-analysis of genome scans and replication identify CD6, IRF8 and TNFRSF1A as new multiple sclerosis susceptibility loci. *Nat. Genet.* 41, 776–782.
- Desmond, D.W., Moroney, J.T., Paik, M.C., Sano, M., Mohr, J.P., Aboumatar, S., Tseng, C.L., Chan, S., Williams, J.B., Remien, R.H., Hauser, W.A., Stern, Y., 2000. Frequency and clinical determinants of dementia after ischemic stroke. *Neurology* 54, 1124–1131.
- Desmond, D.W., Tatemichi, T.K., Paik, M., Stern, Y., 1993. Risk factors for cerebrovascular disease as correlates of cognitive function in a stroke-free cohort. *Arch. Neurol.* 50, 162–166.
- Dong, H., Zitt, C., Auriga, C., Hatzelmann, A., Epstein, P.M., 2010. Inhibition of PDE3, PDE4 and PDE7 potentiates glucocorticoid-induced apoptosis and overcomes glucocorticoid resistance in CEM T leukemic cells. *Biochem. Pharmacol.* 79, 321–329.
- Eichner, J.E., Dunn, S.T., Perveen, G., Thompson, D.M., Stewart, K.E., Stroehla, B.C., 2002. Apolipoprotein E polymorphism and cardiovascular disease: a HuGE review. *Am. J. Epidemiol.* 155, 487–495.
- Evans, D.A., Bennett, D.A., Wilson, R.S., Bienias, J.L., Morris, M.C., Scherr, P.A., Hebert, L.E., Aggarwal, N., Beckett, L.A., Joglekar, R., Berry-Kravis, E., Schneider, J., 2003. Incidence of Alzheimer disease in a biracial urban community: relation to apolipoprotein E allele status. *Arch. Neurol.* 60, 185–189.
- Feskens, E.J., Havekes, L.M., Kalmijn, S., de Knijff, P., Launer, L.J., Kromhout, D., 1994. Apolipoprotein e4 allele and cognitive decline in elderly men. *BMJ* 309, 1202–1206.
- Grodstein, F., Chen, J., Wilson, R.S., Manson, J.E., Study, N.H., 2001. Type 2 diabetes and cognitive function in community-dwelling elderly women. *Diabetes Care* 24, 1060–1065.
- Haan, M.N., Shemanski, L., Jagust, W.J., Manolio, T.A., Kuller, L., 1999. The role of APOE epsilon4 in modulating effects of other risk factors for cognitive decline in elderly persons. *JAMA* 282, 40–46.
- Harold, D., Abraham, R., Hollingworth, P., Sims, R., Gerrish, A., Hamshere, M.L., Pahwa, J.S., Moskva, V., Dowzell, K., Williams, A., Jones, N., Thomas, C., Stretton, A., Morgan, A.R., Lovestone, S., Powell, J., Proitsi, P., Lupton, M.K., Brayne, C., Rubinsztein, D.C., Gill, M., Lawlor, B., Lynch, A., Morgan, K., Brown, K.S., Passmore, P.A., Craig, D., McGuinness, B., Todd, S., Holmes, C., Mann, D., Smith, A.D., Love, S., Kehoe, P.G., Hardy, J., Mead, S., Fox, N., Rossor, M., Collinge, J., Maier, W., Jessen, F., Schürmann, B., van den Bussche, H., Heuser, I., Kornhuber, J., Wiltfang, J., Dichgans, M., Frölich, L., Hampel, H., Hüll, M., Rujescu, D., Goate, A.M., Kauwe, J.S.K., Cruchaga, C., Nowotny, P., Morris, J.C., Mayo, K., Sleegers, K., Bettens, K., Engelborghs, S., De Deyn, P.P., van Broeckhoven, C., Livingston, G., Bass, N.J., Gurling, H., McQuillin, A., Gwilliam, R., Deloukas, P., Al-Chalabi, A., Shaw, C.E., Tzolaki, M., Singleton, A.B., Guerreiro, R., Mühleisen, T.W., Nöthen, M.M., Moebus, S., Jöckel, K.-H., Klopp, N., Wichmann, H.-E., Carrasquillo, M.M., Pankratz, V.S., Younkin, S.G., Holmans, P.A., O'Donovan, M., Owen, M.J., Williams, J., 2009. Genome-wide association study identifies variants at CLU and PICALM associated with Alzheimer's disease. *Nat. Genet.* 41, 1088–1093.
- Henderson, A.S., Eastel, S., Jorm, A.F., Mackinnon, A.J., Korten, A.E., Christensen, H., Croft, L., Jacomb, P.A., 1995. Apolipoprotein E allele epsilon 4, dementia, and cognitive decline in a population sample. *Lancet* 346, 1387–1390.
- Hollingworth, P., Harold, D., Sims, R., Gerrish, A., Lambert, J.-C., Carrasquillo, M.M., Abraham, R., Hamshere, M.L., Pahwa, J.S., Moskva, V., Dowzell, K., Jones, N., Stretton, A., Thomas, C., Richards, A., Ivanov, D., Widdowson, C., Chapman, J., Lovestone, S., Powell, J., Proitsi, P., Lupton, M.K., Brayne, C., Rubinsztein, D.C., Gill, M., Lawlor, B., Lynch, A., Brown, K.S., Passmore, P.A., Craig, D., McGuinness, B., Todd, S., Holmes, C., Mann, D., Smith, A.D., Beaumont, H., Warden, D., Wilcock, G., Love, S., Kehoe, P.G., Hooper, N.M., Vardy, E.R.L.C., Hardy, J., Mead, S., Fox, N.C., Rossor, M., Collinge, J., Maier, W., Jessen, F., Rütger, E., Schürmann, B., Heun, R., Kölsch, H., van den Bussche, H., Heuser, I., Kornhuber, J., Wiltfang, J., Dichgans, M., Frölich, L., Hampel, H., Gallacher, J., Hüll, M., Rujescu, D., Giegling, I., Goate, A.M., Kauwe, J.S.K., Cruchaga, C., Nowotny, P., Morris, J.C., Mayo, K., Sleegers, K., Bettens, K., Engelborghs, S., De Deyn, P.P., Van Broeckhoven, C., Livingston, G., Bass, N.J., Gurling, H., McQuillin, A., Gwilliam, R., Deloukas, P., Al-Chalabi, A., Shaw, C.E., Tzolaki, M., Singleton, A.B., Guerreiro, R., Mühleisen, T.W., Nöthen, M.M., Moebus, S., Jöckel, K.-H., Klopp, N., Wichmann, H.-E., Pankratz, V.S., Sando, S.B., Aasly, J.O., Barcikowska, M., Wszolek, Z.K., Dickson, D.W., Graff-Radford, N.R., Petersen, R.C., van Duijn, C.M., Breteler, M.M.B., Ikram, M.A., DeStefano, A.L., Fitzpatrick, A.L., Lopez, O., Launer, L.J., Seshadri, S., Berr, C., Campion, D., Epelbaum, J., Dartigues, J.-F., Tzourio, C., Alperovitch, A., Lathrop, M., Feulner, T.M., Friedrich, P., Riehle, C., Krawczak, M., Schreiber, S., Mayhaus, M., Nicolhaus, S., Wagenpfeil, S., Steinberg, S., Stefansson, H., Stefansson, K., Snædal, J., Björnsson, S., Jonsson, P.V., Chouraki, V., Genier-Boley, B., Hiltunen, M., Soininen, H., Combarros, O., Zelenika, D., Delepine, M., Bullido, M.J., Pasquier, F., Mateo, I., Frank-Garcia, A., Porcellini, E., Hanon, O., Coto, E., Alvarez, V., Bosco, P., Siciliano, G., Mancuso, M., Panza, F., Solfrizzi, V., Nacmias, B., Sorbi, S., Bossù, P., Piccardi, P., Arosio, B., Annoni, G., Seripa, D., Pilotto, A., Scarpini, E., Galimberti, D., Brice, A., Hannequin, D., Licastro, F., Jones, L., Holmans, P.A., Jonsson, T., Riemenschneider, M., Morgan, K., Younkin, S.G., Owen, M.J., O'Donovan, M., Amouyel, P., Williams, J., 2011. Common variants at



- ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat. Genet.* 43, 429–435.
- Hyman, B.T., Gomez-Isla, T., Briggs, M., Chung, H., Nichols, S., Kohout, F., Wallace, R., 1996. Apolipoprotein E and cognitive change in an elderly population. *Ann. Neurol.* 40, 55–66.
- International HapMap Consortium, Frazer, K.A., Ballinger, D.G., Cox, D.R., Hinds, D.A., Stuve, L.L., Gibbs, R.A., Belmont, J.W., Boudreau, A., Hardenbol, P., Leal, S.M., Pasternak, S.M., Wheeler, D.A., Willis, T.D., Yu, F., Yang, H., Zeng, C., Gao, Y., Hu, H., Hu, H., Li, C., Lin, W., Liu, S., Pan, H., Tang, X., Wang, J., Wang, J., Yu, J., Zhang, B., Zhang, B., Zhao, H., Zhou, J., Gabriel, S.B., Barry, R., Blumenstiel, B., Camargo, A., Defelice, M., Faggart, M., Defelice, M., Gupta, S., Moore, J., Nguyen, H., Onofrio, R.C., Parkin, M., Roy, J., Stahl, E., Winchester, E., Ziaugra, L., Althuler, D., Shen, Y., Yao, Z., Huang, W., Chu, X., He, Y., Jin, L., Liu, Y., Sun, W., Wang, H., Wang, Y., Xiong, X., Xu, L., Waye, M.M., Tsui, S.K., Xue, H., Wong, J.T., Galver, L.M., Fan, J.B., Gunderson, K., Murray, S.S., Oliphant, A.R., Chee, M.S., Montpetit, A., Chagnon, F., Ferretti, V., Leboeuf, M., Olivier, J.F., Phillips, M.S., Roumy, S., Sallee, C., Verner, A., Hudson, T.J., Kwok, P.Y., Cai, D., Koboldt, D.C., Miller, R.D., Pawlikowska, L., Taillon-Miller, P., Xiao, M., Tsui, L.C., Mak, W., Song, Y.Q., Tam, P.K., Nakamura, Y., Kawaguchi, T., Kitamoto, T., Morizono, T., Nagashima, A., Ohnishi, Y., Sekine, A., Tanaka, T., Tsunoda, T., Deloukas, P., Bird, C.P., Delgado, M., Dermitzakis, E.T., Gwilliam, R., Hunt, S., Morrison, J., Powell, D., Stranger, B.E., Whittaker, P., Bentley, D.R., Daly, M.J., de Bakker, P.I., Barrett, J., Chretien, Y.R., Maller, J., McCarroll, S., Patterson, N., Pe'er, I., Price, A., Purcell, S., Richter, D.J., Sabeti, P., Saxena, R., Schaffner, S.F., Sham, P.C., Vairily, P., Stein, L.D., Krishnan, L., Smith, A.V., Tello-Ruiz, M.K., Thorisson, G.A., Chakravarti, A., Chen, P.E., Cutler, D.J., Kashuk, C.S., Lin, S., Abecasis, G.R., Guan, W., Li, Y., Munro, H.M., Qin, Z.S., Thomas, D.J., McVean, G., Auton, A., Bottolo, L., Cardin, N., Eyheramendy, S., Freeman, C., Marchini, J., Myers, S., Spencer, C., Stephens, M., Donnelly, P., Cardon, L.R., Clarke, G., Evans, D.M., Morris, A.P., Weir, B.S., Mullikin, J.C., Sherry, S.T., Feolo, M., Skol, A., Zhang, H., Matsuda, I., Fukushima, Y., Macer, D.R., Suda, E., Rotimi, C.N., Adebamowo, C.A., Ajayi, I., Aniagwu, T., Marshall, P.A., Nkwdimamah, C., Royal, C.D., Leppert, M.F., Dixon, M., Peiffer, A., Qiu, R., Kent, A., Kato, K., Niihara, N., Adewole, I.F., Knoppers, B.M., Foster, M.W., Clayton, E.W., Watkin, J., Muzny, D., Nazareth, L., Sodergren, E., Weinstock, G.M., Yakub, I., Birren, B.W., Wilson, R.K., Fulton, L.L., Rogers, J., Burton, J., Carter, N.P., Clee, C.M., Griffiths, M., Jones, M.C., McLay, K., Plumb, R.W., Ross, M.T., Sims, S.K., Willey, D.L., Chen, Z., Han, H., Kang, L., Godbout, M., Wallenburg, J.C., L'Archeveque, P., Bellemare, G., Saeki, K., An, D., Fu, H., Li, Q., Wang, Z., Wang, R., Holden, A.L., Brooks, L.D., McEwen, J.E., Guyer, M.S., Wang, V.O., Peterson, J.L., Shi, M., Spiegel, J., Sung, L.M., Zacharia, L.F., Collins, F.S., Kennedy, K., Jamieson, R., Stewart, J., Fu, H., Li, Q., Wang, Z., Wang, R., Holden, A.L., Brooks, L.D., McEwen, J.E., Guyer, M.S., Wang, V.O., Peterson, J.L., Shi, M., Spiegel, J., Sung, L.M., Zacharia, L.F., Collins, F.S., Kennedy, K., Jamieson, R., Stewart, J., 2007. A second generation human haplotype map of over 3.1 million SNPs. *Nature* 449, 851–861.
- Israely, I., Costa, R.M., Xie, C.W., Silva, A.J., Kosik, K.S., Liu, X., 2004. Deletion of the neuron-specific protein delta-catenin leads to severe cognitive and synaptic dysfunction. *Curr. Biol.* 14, 1657–1663.
- Johnson, A.D., Handsaker, R.E., Pulit, S.L., Nizzari, M.M., O'Donnell, C.J., de Bakker, P.I., 2008. SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* 24, 2938–2939.
- Jonker, C., Schmand, B., Lindeboom, J., Havekes, L.M., Launer, L.J., 1998. Association between apolipoprotein E epsilon4 and the rate of cognitive decline in community-dwelling elderly individuals with and without dementia. *Arch. Neurol.* 55, 1065–1069.
- Kadoshima-Yamaoka, K., Murakawa, M., Goto, M., Tanaka, Y., Inoue, H., Murafuji, H., Nagahira, A., Hayashi, Y., Nagahira, K., Miura, K., Nakatsuka, T., Chamoto, K., Fukuda, Y., Nishimura, T., 2009. ASB16165, a novel inhibitor for phosphodiesterase 7A (PDE7A), suppresses IL-12-induced IFN-gamma production by mouse activated T lymphocytes. *Immunol. Lett.* 122, 193–197.
- Kim, J.S., Han, S.R., Chung, S.W., Kim, B.S., Lee, K.S., Kim, Y.I., Yang, D.W., Kim, K.S., Kim, J.W., 2003. The apolipoprotein E epsilon4 haplotype is an important predictor for recurrence in ischemic cerebrovascular disease. *J. Neurol. Sci.* 206, 31–37.
- Laird, N.M., Ware, J.H., 1982. Random-effects models for longitudinal data. *Biometrics* 38, 963–974.
- Lambert, J.C., Heath, S., Even, G., Campion, D., Sleegers, K., Hiltunen, M., Combarros, O., Zelenika, D., Bullido, M.J., Tavernier, B., Letenneur, L., Bettens, K., Berr, C., Pasquier, F., Fiévet, N., Barberger-Gateau, P., Engelborghs, S., De Deyn, P., Mateo, I., Franck, A., Helisalmi, S., Porcellini, E., Hanon, O., European Alzheimer's Disease Initiative Investigators, de Pancorbo, M.M., Lendon, C., Dufouil, C., Jaillard, C., Leveillard, T., Alvarez, V., Bosco, P., Mancuso, M., Panza, F., Nacmias, B., Bossù, P., Piccardi, P., Annoni, G., Seripa, D., Galimberti, D., Hannequin, D., Licastro, F., Soininen, H., Ritchie, K., Blanché, H., Dartigues, J.F., Tzourio, C., Gut, I., Van Broeckhoven, C., Alperovitch, A., Lathrop, M., Amouyel, P., 2009. Genome-wide association study identifies variants at CLU and CR1 associated with Alzheimer's disease. *Nat. Genet.* 41, 1094–1099.
- Launer, L.J., Petrovitch, H., Ross, G.W., Markesbery, W., White, L.R., 2008. AD brain pathology: vascular origins? Results from the HAAS autopsy study. *Neurobiol. Aging* 29, 1587–1590.
- Lucin, K.M., Wyss-Coray, T., 2009. Immune activation in brain aging and neurodegeneration: too much or too little? *Neuron* 64, 110–122.
- Martinez-Vicente, M., Cuervo, A.M., 2007. Autophagy and neurodegeneration: when the cleaning crew goes on strike. *Lancet Neurol.* 6, 352–361.
- McCarron, M.O., Delong, D., Alberts, M.J., 1999. APOE genotype as a risk factor for ischemic cerebrovascular disease: a meta-analysis. *Neurology* 53, 1308–1311.
- McClearn, G.E., Johansson, B., Berg, S., Pedersen, N.L., Ahern, F., Pettrill, S.A., Plomin, R., 1997. Substantial genetic influence on cognitive abilities in twins 80 or more years old. *Science* 276, 1560–1563.
- McKhann, G., Drachman, D., Folstein, M., Katzman, R., Price, D., Stadlan, E.M., 1984. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 34, 939–944.
- McQueen, M.B., Bertram, L., Lange, C., Becker, K.D., Albert, M.S., Tanzi, R.E., Blacker, D., 2007. Exploring candidate gene associations with neuropsychological performance. *Am. J. Med. Genet. B Neuro-psychiatr. Genet.* 144B, 987–991.
- Medina, M., Marinescu, R.C., Overhauser, J., Kosik, K.S., 2000. Hemizygosity of delta-catenin (CTNND2) is associated with severe mental retardation in cri-du-chat syndrome. *Genomics* 63, 157–164.
- Miró, X., Pérez-Torres, S., Palacios, J.M., Puigdomènech, P., Mengod, G., 2001. Differential distribution of cAMP-specific phosphodiesterase 7A mRNA in rat brain and peripheral organs. *Synapse* 40, 201–214.
- Monticone, M., Tonachini, L., Tavella, S., Degan, P., Biticchi, R., Palombi, F., Puglisi, R., Boitani, C., Cancedda, R., Castagnola, P., 2007. Impaired expression of genes coding for reactive oxygen species scavenging enzymes in testes of Mtrf1/Chppr-deficient mice. *Reproduction* 134, 483–492.
- Myers, A.J., Gibbs, J.R., Webster, J.A., Rohrer, K., Zhao, A., Marlowe, L., Kaleem, M., Leung, D., Bryden, L., Nath, P., Zismann, V.L., Joshipura, K., Huentelman, M.J., Hu-Lince, D., Coon, K.D., Craig, D.W., Pearson, J.V., Holmans, P., Heward, C.B., Reiman, E.M., Stephan, D., Hardy, J., 2007. A survey of genetic human cortical gene expression. *Nat. Genet.* 39, 1494–1499.
- Naj, A.C., Jun, G., Beecham, G.W., Wang, L.S., Vardarajan, B.N., Buross, J., Gallins, P.J., Buxbaum, J.D., Jarvik, G.P., Crane, P.K., Larson, E.B., Bird, T.D., Boeve, B.F., Graff-Radford, N.R., De Jager, P.L., Evans,

- D., Schneider, J.A., Carrasquillo, M.M., Ertekin-Taner, N., Younkin, S.G., Cruchaga, C., Kauwe, J.S., Nowotny, P., Kramer, P., Hardy, J., Huentelman, M.J., Myers, A.J., Barmada, M.M., Demirci, F.Y., Baldwin, C.T., Green, R.C., Rogava, E., St George-Hyslop, P., Arnold, S.E., Barber, R., Beach, T., Bigio, E.H., Bowen, J.D., Boxer, A., Burke, J.R., Cairns, N.J., Carlson, C.S., Carney, R.M., Carroll, S.L., Chui, H.C., Clark, D.G., Comeveaux, J., Cotman, C.W., Cummings, J.L., DeCarli, C., Dekosky, S.T., Diaz-Arrastia, R., Dick, M., Dickson, D.W., Ellis, W.G., Faber, K.M., Fallon, K.B., Farlow, M.R., Ferris, S., Frosch, M.P., Galasko, D.R., Ganguli, M., Gearing, M., Geschwind, D.H., Ghetti, B., Gilbert, J.R., Gilman, S., Giordani, B., Glass, J.D., Growdon, J.H., Hamilton, R.L., Harrell, L.E., Head, E., Honig, L.S., Hulette, C.M., Hyman, B.T., Jicha, G.A., Jin, L.-W., Johnson, N., Karlawish, J., Karydas, A., Kaye, J.A., Kim, R., Koo, E.H., Kowall, N.W., Lah, J.J., Levey, A.I., Lieberman, A.P., Lopez, O.L., Mack, W.J., Marson, D.C., Martiniuk, F., Mash, D.C., Masliah, E., McCormick, W.C., Mccurry, S.M., McDavid, A.N., McKee, A.C., Mesulam, M., Miller, B.L., Miller, C.A., Miller, J.W., Parisi, J.E., Perl, D.P., Peskind, E., Petersen, R.C., Poon, W.W., Quinn, J.F., Rajbhandary, R.A., Raskind, M., Reisberg, B., Ringman, J.M., Roberson, E.D., Rosenberg, R.N., Sano, M., Schneider, L.S., Seeley, W., Sheltanski, M.L., Slifer, M.A., Smith, C.D., Sonnen, J.A., Spina, S., Stern, R.A., Tanzi, R.E., Trojanowski, J.Q., Troncoso, J.C., Deerlin, V.M.V., Vinters, H.V., Vonsattel, J.P., Weintraub, S., Welsh-Bohmer, K.A., Williamson, J., Woltjer, R.L., Cantwell, L.B., Dombroski, B.A., Beekly, D., Lunetta, K.L., Martin, E.R., Kamboh, M.I., Saykin, A.J., Reiman, E.M., Bennett, D.A., Morris, J.C., Montine, T.J., Goate, A.M., Blacker, D., Tsuang, D.W., Hakonarson, H., Kukull, W.A., Foroud, T.M., Haines, J.L., Mayeux, R., Pericak-Vance, M.A., Farrer, L.A., Schellenberg, G.D., 2011. Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nat. Genet.* 43, 436–441.
- Neuropathology Group of the Medical Research Council Cognitive Function and Ageing Study, 2001. Pathological correlates of late-onset dementia in a multicentre, community-based population in England and Wales. Neuropathology Group of the Medical Research Council Cognitive Function and Ageing Study (MRC CFAS). *Lancet* 357, 169–175.
- Park, J.H., Wacholder, S., Gail, M.H., Peters, U., Jacobs, K.B., Chanock, S.J., Chatterjee, N., 2010. Estimation of effect size distribution from genome-wide association studies and implications for future discoveries. *Nat. Genet.* 42, 570–575.
- Pendlebury, S.T., Rothwell, P.M., 2009. Prevalence, incidence, and factors associated with pre-stroke and post-stroke dementia: a systematic review and meta-analysis. *Lancet Neurol.* 8, 1006–1018.
- Pérez-Torres, S., Cortés, R., Tolnay, M., Probst, A., Palacios, J.M., Mengod, G., 2003. Alterations on phosphodiesterase type 7 and 8 isozyme mRNA expression in Alzheimer's disease brains examined by in situ hybridization. *Exp. Neurol.* 182, 322–334.
- Price, A.L., Patterson, N.J., Plenge, R.M., Weinblatt, M.E., Shadick, N.A., Reich, D., 2006. Principal components analysis corrects for stratification in genome-wide association studies. *Nat. Genet.* 38, 904–909.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., de Bakker, P.I., Daly, M.J., Sham, P.C., 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81, 559–575.
- Reijmer, Y.D., van den Berg, E., Ruis, C., Kappelle, L.J., Biessels, G.J., 2010. Cognitive dysfunction in patients with type 2 diabetes. *Diabetes/Metab. Res. Rev.* 26, 507–519.
- Reitz, C., Brayne, C., Mayeux, R., 2011a. Epidemiology of Alzheimer Disease. *Nat. Rev. Neurol.* 7, 137–152.
- Reitz, C., Tokunishi, S., Clark, L.N., Conrad, C., Vonsattel, J.-P., Hazrati, L.-N., Palotás, A., Lantigua, R., Medrano, M., Jiménez-Velázquez, I., Vardarajan, B., Simkin, I., Haines, J.L., Pericak-Vance, M.A., Farrer, L.A., Lee, J.H., Rogava, E., George-Hyslop, P.S., Mayeux, R., 2011b. SORCS1 alters amyloid precursor protein processing and variants may increase Alzheimer's disease risk. *Ann. Neurol.* 69, 47–64.
- Rhead, B., Karolchik, D., Kuhn, R.M., Hinrichs, A.S., Zweig, A.S., Fujita, P.A., Diekhans, M., Smith, K.E., Rosenbloom, K.R., Raney, B.J., Pohl, A., Pheasant, M., Meyer, L.R., Learned, K., Hsu, F., Hillman-Jackson, J., Harte, R.A., Giardine, B., Dreszer, T.R., Clawson, H., Barber, G.P., Haussler, D., Kent, W.J., 2010. The UCSC Genome Browser database: update 2010. *Nucleic Acids Res.* 38, D613–D619.
- Ross, C.A., Poirier, M.A., 2004. Protein aggregation and neurodegenerative disease. *Nat. Med.* 10, S10–S17.
- Rowe, J.A., Moulds, J.M., Newbold, C.I., Miller, L.H., 1997. P. falciparum rosetting mediated by a parasite-variant erythrocyte membrane protein and complement-receptor 1. *Nature* 388, 292–295.
- Schneider, J.A., Arvanitakis, Z., Bang, W., Bennett, D.A., 2007. Mixed brain pathologies account for most dementia cases in community-dwelling older persons. *Neurology* 69, 2197–2204.
- Schneider, J.A., Bennett, D.A., 2010. Where vascular meets neurodegenerative disease. *Stroke* 41 (Suppl e), S144–S156.
- Scott, L.J., Mohlke, K.L., Bonnycastle, L.L., Willer, C.J., Li, Y., Duren, W.L., Erdos, M.R., Stringham, H.M., Chines, P.S., Jackson, A.U., Prokunina-Olsson, L., Ding, C.J., Swift, A.J., Narisu, N., Hu, T., Pruim, R., Xiao, R., Li, X.Y., Conneely, K.N., Riebow, N.L., Sprau, A.G., Tong, M., White, P.P., Hetrick, K.N., Barnhart, M.W., Bark, C.W., Goldstein, J.L., Watkins, L., Xiang, F., Saramies, J., Buchanan, T.A., Watanabe, R.M., Valle, T.T., Kinnunen, L., Abecasis, G.R., Pugh, E.W., Doheny, K.F., Bergman, R.N., Tuomilehto, J., Collins, F.S., Boehnke, M., 2007. A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science* 316, 1341–1345.
- Seshadri, S., Fitzpatrick, A.L., Ikram, M.A., DeStefano, A.L., Gudnason, V., Boada, M., Bis, J.C., Smith, A.V., Carrasquillo, M.M., Lambert, J.C., Harold, D., Schrijvers, E.M.C., Ramirez-Lorca, R., Debette, S., Longstreth, W.T., Janssens, A.C.J.W., Pankratz, V.S., Dartigues, J.F., Hollingworth, P., Aspeland, T., Hernandez, I., Beiser, A., Kuller, L.H., Koudstaal, P.J., Dickson, D.W., Tzourio, C., Abraham, R., Antunez, C., Du, Y., Rotter, J.I., Aulchenko, Y.S., Harris, T.B., Petersen, R.C., Berr, C., Owen, M.J., Lopez-Arrieta, J., Varadarajan, B.N., Becker, J.T., Rivadeneira, F., Nalls, M.A., Graff-Radford, N.R., Campion, D., Auerbach, S., Rice, K., Hofman, A., Jonsson, P.V., Schmidt, H., Lathrop, M., Mosley, T.H., Au, R., Psaty, B.M., Uitterlinden, A.G., Farrer, L.A., Lumley, T., Ruiz, A., Williams, J., Amouyel, P., Younkin, S.G., Wolf, P.A., Launer, L.J., Lopez, O.L., van Duijn, C.M., Breteler, M.M.B., Consortium, C., 2010. Genome-wide analysis of genetic loci associated with Alzheimer disease. *JAMA* 303, 1832–1840.
- Sonnen, J.A., Larson, E.B., Crane, P.K., Haneuse, S., Li, G., Schellenberg, G.D., Craft, S., Leverenz, J.B., Montine, T.J., 2007. Pathological correlates of dementia in a longitudinal, population-based sample of aging. *Ann. Neurol.* 62, 406–413.
- Stern, Y., 2009. Cognitive reserve. *Neuropsychologia* 47, 2015–2028.
- Stranger, B.E., Nica, A.C., Forrester, M.S., Dimas, A., Bird, C.P., Beazley, C., Ingle, C.E., Dunning, M., Flicek, P., Koller, D., Montgomery, S., Tavaré, S., Deloukas, P., Dermitzakis, E.T., 2007. Population genomics of human gene expression. *Nat. Genet.* 39, 1217–1224.
- Su, A.I., Cooke, M.P., Ching, K.A., Hakak, Y., Walker, J.R., Wiltshire, T., Orth, A.P., Vega, R.G., Sapinoso, L.M., Moqrich, A., Patapoutian, A., Hampton, G.M., Schultz, P.G., Hogenesch, J.B., 2002. Large-scale analysis of the human and mouse transcriptomes. *Proc. Natl. Acad. Sci. U. S. A.* 99, 4465–4470.
- Swan, G.E., Carmelli, D., Reed, T., Harshfield, G.A., Fabsitz, R.R., Eslinger, P.J., 1990. Heritability of cognitive performance in aging twins. The National Heart, Lung, and Blood Institute Twin Study. *Arch. Neurol.* 47, 259–262.
- Tonachini, L., Monticone, M., Puri, C., Tacchetti, C., Pinton, P., Rizzuto, R., Cancedda, R., Tavella, S., Castagnola, P., 2004. Chondrocyte protein with a poly-proline region (CHPPR) is a novel mitochondrial

- protein and promotes mitochondrial fission. *J. Cell. Physiol.* 201, 470–482.
- Troncoso, J.C., Zonderman, A.B., Resnick, S.M., Crain, B., Pletnikova, O., O'Brien, R.J., 2008. Effect of infarcts on dementia in the Baltimore longitudinal study of aging. *Ann. Neurol.* 64, 168–176.
- Warsch, J.R., Wright, C.B., 2010. The aging mind: vascular health in normal cognitive aging. *J. Am. Geriatr. Soc.* 58 (Suppl 2), S319–S324.
- Willer, C.J., Li, Y., Abecasis, G.R., 2010. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26, 2190–2191.
- Wilson, R., Bienias, J., Berry-Kravis, E., Evans, D., Bennett, D., 2002a. The apolipoprotein E epsilon 2 allele and decline in episodic memory. *J. Neurol. Neurosurg., Psychiatry* 73, 672–677.
- Wilson, R., Schneider, J., Barnes, L., Beckett, L., Aggarwal, N., Cochran, E., Berry-Kravis, E., Bach, J., Fox, J., Evans, D., Bennett, D., 2002b. The apolipoprotein E epsilon 4 allele and decline in different cognitive systems during a 6-year period. *Arch. Neurol.* 59, 1154–1160.
- Wilson, R.S., Barnes, L.L., Krueger, K.R., Hoganson, G., Bienias, J.L., Bennett, D.A., 2005. Early and late life cognitive activity and cognitive systems in old age. *J. Int. Neuropsychol. Soc.* 11, 400–407.
- Wilson, R.S., Beckett, L.A., Barnes, L.L., Schneider, J.A., Bach, J., Evans, D.A., Bennett, D.A., 2002c. Individual differences in rates of change in cognitive abilities of older persons. *Psychol. Aging* 17, 179–193.
- Wilson, R.S., Gilley, D.W., Bennett, D.A., Beckett, L.A., Evans, D.A., 2000. Person-specific paths of cognitive decline in Alzheimer's disease and their relation to age. *Psychol. Aging* 15, 18–28.
- Wilson, R.S., Leurgans, S.E., Boyle, P.A., Schneider, J.A., Bennett, D.A., 2010. Neurodegenerative basis of age-related cognitive decline. *Neurology* 75, 1070–1078.
- Wu, C., Orozco, C., Boyer, J., Leglise, M., Goodale, J., Batalov, S., Hodge, C.L., Haase, J., Janes, J., Huss, J.W., Su, A.I., 2009. BioGPS: an extensible and customizable portal for querying and organizing gene annotation resources. *Genome Biol.* 10, R130.
- Yang, G., McIntyre, K.W., Townsend, R.M., Shen, H.H., Pitts, W.J., Dodd, J.H., Nadler, S.G., McKinnon, M., Watson, A.J., 2003. Phosphodiesterase 7A-deficient mice have functional T cells. *J. Immunol.* 171, 6414–6420.