

# Detecting genetic association through shortest paths in a bidirected graph

Masao Ueki<sup>1</sup> | Yoshinori Kawasaki<sup>2</sup> | Gen Tamiya<sup>3</sup> | for Alzheimer's Disease  
Neuroimaging Initiative

<sup>1</sup>Biostatistics Center, Kurume University, Fukuoka, Japan

<sup>2</sup>Department of Statistical Modeling, The Institute of Statistical Mathematics, The Graduate University for Advanced Studies, Tachikawa, Tokyo, Japan

<sup>3</sup>Statistical Genetics and Genomics, Tohoku Medical Megabank Organization, Tohoku University, Aoba-Ku, Sendai, Japan

## Correspondence

Masao Ueki, Biostatistics Center, Kurume University, 67 Asahi-Machi, Kurume, Fukuoka 830-0011, Japan.

Email: uekimrsd@nifty.com

## ABSTRACT

Genome-wide association studies (GWASs) commonly use marginal association tests for each single-nucleotide polymorphism (SNP). Because these tests treat SNPs as independent, their power will be suboptimal for detecting SNPs hidden by linkage disequilibrium (LD). One way to improve power is to use a multiple regression model. However, the large number of SNPs preclude simultaneous fitting with multiple regression, and subset regression is infeasible because of an exorbitant number of candidate subsets. We therefore propose a new method for detecting hidden SNPs having significant yet weak marginal association in a multiple regression model. Our method begins by constructing a bidirected graph locally around each SNP that demonstrates a moderately sized marginal association signal, the *focal* SNPs. Vertexes correspond to SNPs, and adjacency between vertexes is defined by an LD measure. Subsequently, the method collects from each graph all shortest paths to the focal SNP. Finally, for each shortest path the method fits a multiple regression model to all the SNPs lying in the path and tests the significance of the regression coefficient corresponding to the terminal SNP in the path. Simulation studies show that the proposed method can detect susceptibility SNPs hidden by LD that go undetected with marginal association testing or with existing multivariate methods. When applied to real GWAS data from the Alzheimer's Disease Neuroimaging Initiative (ADNI), our method detected two groups of SNPs: one in a region containing the *apolipoprotein E (APOE)* gene, and another in a region close to the *semaphorin 5A (SEMA5A)* gene.

## KEYWORDS

bidirected graph, conservative multiple test, hidden association, linkage disequilibrium, shortest path

Grant sponsor: Alzheimer's Disease Neuroimaging Initiative (ADNI); Grant sponsor: National Institutes of Health; Grant number: U01 AG024904; Grant sponsor: DODADNI; Grant sponsor: National Institute on Aging; Grant sponsor: National Institute of Biomedical Imaging and Bioengineering; Grant sponsor: AbbVie, Alzheimer's Association; Grant sponsor: Alzheimer's Drug Discovery Foundation; Grant sponsor: Araclon Biotech; Grant sponsor: BioClinica, Inc.; Grant sponsor: Biogen; Bristol-Myers Squibb Company; Grant sponsor: CereSpir, Inc.; Grant sponsor: Eisai Inc.; Grant sponsor: Elan Pharmaceuticals, Inc.; Grant sponsor: Eli Lilly and Company; Grant sponsor:

EuroImmun; Grant sponsor: F. Hoffmann-La Roche Ltd.; Grant sponsor: Genentech, Inc.; Grant sponsor: Fujirebio; Grant sponsor: GE Healthcare; Grant sponsor: IXICO Ltd.; Grant sponsor: Janssen Alzheimer Immunotherapy Research & Development, LLC; Grant sponsor: Johnson & Johnson Pharmaceutical Research & Development LLC; Grant sponsor: Lumosity; Grant sponsor: Lundbeck; Grant sponsor: Merck & Co., Inc.; Grant sponsor: Meso Scale Diagnostics, LLC; Grant sponsor: NeuroRx Research; Grant sponsor: Neurotrack Technologies; Grant sponsor: Novartis Pharmaceuticals Corporation; Grant sponsor: Pfizer, Inc.; Grant sponsor: Piramal Imaging; Grant

## 1 | INTRODUCTION

In genome-wide association studies (GWASs), researchers routinely use univariate regression to analyze each single-nucleotide polymorphism (SNP). Although GWASs have led to the discovery of many new SNPs that are involved in disease development and/or progression of clinical characteristics, the genetic architecture of many human traits remains largely unknown (Manolio, 2013). This is referred to as a missing heritability problem (Manolio et al., 2009; Maher, 2008), for which several explanations have been proposed. These include gene-environment interaction, gene-gene interaction, and the contributions of rarer variants, structural variants, and many variants with small effects. Researchers are attempting to develop statistical methods to test such challenging scientific hypotheses. Linkage disequilibrium (LD) may also contribute to missing heritability (Ehret et al., 2012). The univariate regression approach—or marginal association testing—works for SNP panels designed to capture causal variants under the assumption of a single causal variant or unlinked multiple causal variants (de Bakker et al., 2005; Eberle et al., 2007; Spencer, Su, Donnelly, & Marchini, 2009). If this assumption is violated, marginal association testing, which treats SNPs as independent and does not fully take LD into account, may fail to achieve adequate statistical power. For example, if multiple causal SNPs in a region are in LD, the marginal association signal can be weakened due to cancelation of effects (Eberle et al., 2007; for an illustration under a bivariate model, see supplementary information Section 4). We refer to this as the *hidden signal* problem. It differs from the case of an untyped causal variant, where the true causal variant is not included in the panel of SNPs tested. The hidden signal problem can occur even if all causal variants are included in the data. Therefore, an association signal hidden by LD may be undetectable, even with genotype imputation, when a marginal association test is employed.

---

sponsor: Servier; Grant sponsor: Takeda Pharmaceutical Company; Grant sponsor: Transition Therapeutics; Grant sponsor: Canadian Institutes of Health Research; Grant sponsor: Foundation for the National Institutes of Health ([www.fnih.org](http://www.fnih.org)); Grant sponsor: Northern California Institute for Research and Education; Grant sponsor: Laboratory for Neuro Imaging at the University of Southern California; Grant number: 2015-ISM-CRP-1013; Grant sponsor: JSPS KAKENHI; Grant numbers: JP16K00064, JP16K08638, JP16H05242, JP15H01678, JP16H01528

Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database ([adni.loni.usc.edu](http://adni.loni.usc.edu)). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at [http://adni.loni.usc.edu/wp-content/uploads/how\\_to\\_apply/ADNI\\_Acknowledgement\\_List.pdf](http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf)

Simultaneous analysis using multiple regression may be effective if the number of candidate SNPs is small. However, because GWAS data contain a large number of SNPs, fitting them simultaneously with a multiple regression model is impossible. A two-marker test based on bivariate regression is applicable to the above problem: Kim, Morris, Won, and Elston (2010), Slavin, Feng, Schnell, Zhu, and Elston (2011), and Wang, Morris, Schaid, and Elston (2012) proposed using two adjacent SNPs, and Howey and Cordell (2014) recently proposed a different two-marker test, an artificial imputation test. All these two-marker tests are motivated by the goal of improving statistical power under a scenario in which a single causal variant exists and coverage by the marker (tag) SNPs is poor. Another applicable method is the multiple marker test based on a sliding window approach, as implemented in the UNPHASED method (Dudbridge, 2008; Dudbridge, F., Holmans, P. A., & Wilson, 2011); this approach tests the effect of a haplotype comprising multiple SNPs. The sliding window approach tests the combined, group effect of a set of nearby SNPs. However, it is not necessary that nearby SNPs possess similar effects. Taking a wider window may avoid missing causal SNPs, but it concomitantly increases the number of tested SNPs, thereby expending degrees of freedom—in particular, when most of the SNPs in a group are neutral—so that statistical power is greatly decreased.

In this paper, we describe a new method that can detect SNPs hidden by LD. To account for LD, we fit a multiple regression model for each SNP one at a time (which we call the *focal* SNP) by incorporating other, adjacent SNPs that are in LD with the focal SNP. We develop an SNP selection method based on a bidirected (or covariance) graph (Cox & Wermuth, 1996; Drton & Perlman, 2007; Kauermann, 1996) to narrow the field of candidate SNPs to be incorporated into the model. Adjacency is defined using a measure of LD. Then, in the bidirected graph, all shortest paths to the focal SNP are identified and saved. Finally, the multiple regression model is fitted to all of the SNPs lying in each shortest path, and significance of the regression coefficient corresponding to the terminal SNP (the SNP farthest from the focal SNP) in each path is tested. The rationale for seeking the shortest paths is that the regression coefficient of the terminal SNP has nonzero effect in a noiseless setting (see Proposition 2). Our method aims to detect SNPs that have weak marginal association but show strong association in a multiple regression model after adjustment is made for confounding due to multiple nearby SNPs. We develop a conservative multiple testing procedure to control the family-wise error rate in testing the regression coefficients of the terminal SNPs in each graph. Simulation studies and application to real GWAS data from the Alzheimer's Disease Neuroimaging Initiative (ADNI) demonstrate that the proposed method can detect disease susceptibility SNPs hidden by LD that go undetected with the marginal association test or with existing multivariate methods.

## 2 | MATERIALS AND METHODS

### 2.1 | The proposed bidirected graph approach

Suppose that we have observations consisting of  $n$  subjects with  $p$  SNPs  $X = (X_1, \dots, X_p)$ , where  $X_j = (X_{1j}, \dots, X_{nj})^T$  with  $X_{ij} \in \{0, 1, 2\}$  (i.e., the minor allele count) for  $i = 1, \dots, n$  and  $j = 1, \dots, p$ , together with quantitative phenotype data  $y = (y_1, \dots, y_n)^T$ . Let  $M = \{1, \dots, p\}$ . For a subset  $A \subset M$ , we denote by  $X_A$  the subcolumn matrix of  $X$  in which the indexes run through the subset  $A$ . Similar notation is used for vectors throughout. In standard SNP-GWAS analysis, the effect of each SNP is investigated by testing the null hypothesis that the regression coefficient  $\beta_j$  of  $j$ th SNP is 0 in a marginal model,

$$y = 1\beta_0 + X_j\beta_j + \epsilon,$$

where  $\beta_0$  is the intercept,  $1$  denotes an  $n$ -vector with all components being unity, and  $\epsilon \sim N(0, \sigma^2 I)$ . When the marginal effect is weak, the standard marginal testing approach is expected to be underpowered. The existence of multiple causal SNPs that are in LD may weaken the marginal association signal (see supplementary information Section 4). In this situation, controlling for the correlation structure due to LD should improve the detection power. Therefore, we consider testing for the effect of the  $j$ th SNP conditional on a set of other SNPs,  $A \subset M \setminus \{j\}$ . Specifically, we test the null


hypothesis that  $\beta_j$ , the regression coefficient of  $j$ th SNP, is 0 in a multiple linear regression model,

$$y = 1\beta_0 + X_j\beta_j + X_A\beta_A + \epsilon,$$


where  $\beta_A$  is the vector of regression coefficients corresponding to  $X_A$  and  $\epsilon \sim N(0, \sigma^2 I)$ . Let  $\hat{\beta}_{j,A}$  be the regression coefficient estimated by least squares in the above multiple regression model. If  $A = \emptyset$ , the above test reduces to the standard marginal association test. To detect SNPs whose effect is hidden by LD, testing for the effect of  $\hat{\beta}_{j,A}$  with a nonempty set  $A \subset M \setminus \{j\}$  may produce higher power than the marginal association test. However, there exist numerous choices for  $A$ , so with GWAS data, where  $p$  is large, it is difficult to compute  $\hat{\beta}_{j,A}$  for all possible subsets  $A \subset M$ . Even if all combinations of  $A$  are computable, we face a multiple testing burden in evaluating  $\hat{\beta}_{j,A}$ , that is, testing the null hypothesis

$$\forall A \subset M \setminus \{j\} : \beta_{j,A} = 0.$$

It is unclear how to carry out a multiple testing correction. There are  $\binom{p-1}{m}$  possibilities of  $\hat{\beta}_{j,A}$  for  $A$  with  $|A| = m$  for a given  $m < n$ . For example, consider  $p = 500,000$  SNPs. Then, for  $m = 2, 3$ , and  $4$  (i.e., two, three, and four conditioning SNPs), we have roughly  $1.2 \times 10^{11}$ ,  $2 \times 10^{16}$ , and  $3 \times 10^{21}$  possibilities, respectively. If all tests are independent, then the Bonferroni correction is appropriate. However, the independence assumption is violated by the correlation among tests due to LD as well as the repeated tests of the  $j$ th SNP simply by altering the set of conditioning SNPs. Therefore, the naive



**FIGURE 1** Illustration of the proposed shortest path test. The focal SNP,  $X_1$ , shows (weak) marginal association with  $\mu = E(y|X)$  based on a moderate  $P$ -value cutoff (path 0). A bidirected graph around SNP  $X_1$  is constructed with adjacency defined by the magnitude of LD, and all shortest paths to  $X_1$  are collected. In this example, there are five shortest paths numbered 1–5, as shown in the top right. For each of the five shortest paths, the SNP corresponding to the terminal vertex is tested for association in a multiple regression model including all SNPs in the path. The fitted multiple regression model is described for each shortest path. The tested SNP and its regression coefficient are emphasized in bold and red, respectively. Conservative multiple testing is adapted in order to take the number of shortest paths into account



**FIGURE 2** Type I error rates from quantitative trait simulations comparing the proposed method with  $(a_{\min}, \alpha_1) \in \{0.1, 0.3, 0.5, 0.7\} \times \{0.01, 0.001, 0.0001, 0.00001\}$ , the marginal association test, and SnipSnip. Nominal family-wise error rates are  $\alpha = 30\%, 20\%, 10\%$ , and  $5\%$

Bonferroni approach should lead to unrealistically stringent multiple testing correction.


To deal with this challenging problem, we utilize a bidirected graph (Drton & Perlman, 2007) for SNPs with adjacency defined by the pairwise correlation between SNPs. For a given matrix  $X$ , let  $Q_X = I - P_X$ , where  $P_X$  is the projection onto the column space of  $X$ . Then, for an  $n$ -dimensional vector  $x$ ,  $Q_X x$  is the residual from regression of  $x$  onto  $X$ . Hereafter, we assume that  $n$ -dimensional vectors are standardized: specifically, assume that  $x$  satisfies both  $x^T 1 = 0$  and  $\|x\|^2 = 1$ . For two standardized  $n$ -dimensional vectors  $x_1$  and  $x_2$ , the inner product  $x_1^T x_2 = \sum_{i=1}^n x_{1i} x_{2i}$ , equals the sample correlation. Centering of an  $n$ -vector  $X_j$  is obtained by  $x_j = Q_1 X_j = X_j - P_1 X_j$ , then  $x_j^T 1 = 0$  holds. In some circumstances, we have covariates  $Z$ , such as age, gender, and known genetic variants, whose effects on  $y$  need to be removed according to prior scientific knowledge. In such cases, we consider  $x_j = Q_{(1,Z)} X_j$ .

We denote the bidirected graph by  $G = (V, E)$  having a vertex set  $V = \{0\} \cup M$  and an edge set  $E \subset V \times V$ . Hence,  $|V| = 1 + p$ . Let the vertex 0 be the phenotype, that is,  $x_0 =$

$\mu = E(y|X)$ . The remaining vertexes belonging to  $M$  correspond to the  $p$  SNPs. Two arbitrary vertexes  $j$  and  $k$  in  $V$  are adjacent if and only if  $(j, k) \in E$ ; otherwise, they are not adjacent. Here, we say that two vertexes  $j$  and  $k$  ( $j \neq k$ ) in  $V$  are adjacent if

$$x_j^T x_k \neq 0. \tag{1}$$

Adjacent vertexes are said to be spouses, and they are joined by a bidirected edge  $j \leftrightarrow k$ . Therefore, absence of an edge between vertexes  $j$  and  $k$  corresponds to  $x_j^T x_k = 0$ , whereas the presence of an edge corresponds to  $x_j^T x_k \neq 0$ . A path from vertex  $j$  to vertex  $k$  is a sequence of adjacent edges connecting  $j$  and  $k$  for which the corresponding sequence of vertexes contains no repetitions. We refer to the first and the last vertexes as the terminals of the path, and all other vertexes as intermediate vertexes. For two standardized SNP vectors,  $x_j$  and  $x_k$ , coded as 0, 1, 2,  $x_j^T x_k$  coincides with the Wellek-Ziegler correlation coefficient, which measures the magnitude of LD—just as with the Pearson correlation coefficient for haploid data—even if the SNPs are unphased. In fact, the



**FIGURE 3** Type I error rates from binary trait simulations comparing the proposed method with  $(a_{\min}, \alpha_1) \in \{0.1, 0.3, 0.5, 0.7\} \times \{0.01, 0.001, 0.0001, 0.00001\}$ , the marginal association test, and SnipSnip. Nominal family-wise error rates are  $\alpha = 30\%, 20\%, 10\%$ , and  $5\%$

population version of the Wellek-Ziegler correlation coefficient coincides with the population Pearson correlation coefficient for haploid data under Hardy-Weinberg equilibrium (Wellek & Ziegler, 2009). With real GWAS data, a value of adjacency  $x_j^T x_k$  exactly equal to 0 is not practical in finite samples. In practice, we introduce a nonzero cutoff value for judging an adjacency of zero in our bidirected graph.

Our proposed approach is based on the shortest paths to  $\mu$  in a bidirected graph constructed locally for each SNP that shows (weak) association with  $y$  with a moderate  $P$ -value cutoff  $\alpha_1$ —we call these SNPs of interest as *focal* SNPs. Using the shortest paths, we can narrow the candidates for  $A$ , thereby rendering computation feasible and reducing the multiple testing burden. The motivation behind our proposal is given in the Appendix. Centered around each focal SNP, say SNP  $X_1$ , which has  $p_1$  nearby SNPs, we compute  $\binom{p_1}{2}$  Wellek-Ziegler correlation coefficients. Subsequently, we collect all of the shortest paths starting from  $x_1$  in the bidirected graph. (We regard the focal SNPs with moderate marginal signals as being adjacent to  $\mu$ .) An illustration of constructing the

bidirected graph is given in Figure 1. If one of the shortest paths is  $\mu \leftrightarrow X_1 \leftrightarrow X_2 \leftrightarrow \dots \leftrightarrow X_l \leftrightarrow X_{l+1}$ , we test the null hypothesis  $\beta_{l+1} = 0$  for the terminal SNP, with the test based on the multiple regression model,

$$y = 1\beta_0 + X_1\beta_1 + X_2\beta_2 + \dots + X_l\beta_l + X_{l+1}\beta_{l+1} + \epsilon. \quad (2)$$

If  $X_1$  and  $\mu$  are truly adjacent, then the path is the shortest between  $\mu$  and SNP  $X_{l+1}$ . Then, Proposition 2 in the Appendix ensures that  $E(\hat{\beta}_{l+1, \{1, \dots, l\}} | X) \neq 0$ , where  $\hat{\beta}_{l+1, \{1, \dots, l\}}$  is the least-squares estimator in (2). However, the adjacency between  $X_1$  and  $\mu$  is arbitrarily determined by some moderate  $P$ -value cutoff. Furthermore, the above statement only mentions nonzero effects but takes no account of stochastic noise. We overcome these issues by testing hypotheses for the  $\hat{\beta}_{l+1, \{1, \dots, l\}}$ . We employ a conservative testing approach for the terminal SNP in each shortest path as described in Section 2.2. For binary phenotypes, we use the multiple logistic regression model with the Wald test.

**TABLE 1** Causal SNPs for power simulations

	rs9423671	rs9423711	rs2388264	rs11251392	rs17155640
rs9423671		82	70	−42	−19
rs9423711			87	−42	−32
rs2388264				−35	−42
rs11251392					−61
rs17155640					
MAF(%)	32	26	26	38	37
Scenario	$\beta_1^*$	$\beta_2^*$	$\beta_3^*$	$\beta_4^*$	$\beta_5^*$
1			0.20		
2	−0.40		0.50		
3		−0.60	0.70		
4	0.40	−0.70	0.50		
5	−0.20		0.50	0.40	0.30
6	0.50	−0.50	0.50	0.50	0.50

Note. Upper table: Wellek–Ziegler correlation coefficients ( $\times 100$ ) and minor allele frequency (%) for five causal SNPs, rs9423671, rs9423711, rs2388264, rs11251392, and rs17155640, whose indexes in the 1,273 SNPs ordered in base pair position are 457, 476, 492, 533, and 534, respectively. Lower table: Regression models considered in power simulations. Five regressions of the form  $\beta_0 + \sum_{j=1}^5 x_{i,j} \beta_{0,j}$  were used, where the true regression coefficients  $\beta_{0,j}$  for the six scenarios were set as  $\beta_j^*$  multiplied by a constant  $m > 0$ , and empty cells indicate that the true coefficient was 0.

## 2.2 | Algorithm description

Here, we describe the proposed algorithm in detail.

- 1. Initialization** Set a  $P$ -value cutoff  $\alpha_1$  for screening, and a desired family-wise error rate  $\alpha$  (e.g., 5%).
  - 2. Marginal association test** Conduct marginal association tests for all SNPs  $k \in M = \{1, \dots, p\}$  under the marginal model  $\beta_0 + X_k \beta_k$ , which yields  $P$ -values  $P_k$ ,  $k \in M$ . Let  $H = \{k \in M : P_k < \alpha_1\}$  be the set of SNPs of interest, the focal SNPs.
  - 3. Test in bidirected graph** For each  $k \in H$ , set SNP  $k$  as the focal SNP. Denote its nearby SNPs by  $N(k) \subset M$  including the focal SNP (SNP  $k$ ) itself, where nearby SNPs are determined by a 100 kb window (see Section 5) both upstream and downstream of the  $k$ th SNP. Build a bidirected graph where adjacency between two vertexes  $l, m \in N(k)$  is declared if  $\rho_{lm}^2 \in [a_{\min}, a_{\max}]$ . Here  $a_{\min}$  and  $a_{\max}$  ( $a_{\min} < a_{\max}$ ) are pre-specified cutoff values, and  $\rho_{lm}$  is the Wellek–Ziegler correlation coefficient (Pearson correlation coefficient for SNP genotypes coded 0,1,2) between the  $l$ th and  $m$ th SNPs. Denote the resulting bidirected graph by  $G_k$ .
- 3.1. Collection of shortest paths to centered SNP  $k$**  Collect all shortest paths from the focal SNP  $k$  to each vertex in  $G_k$  and denote the set of all shortest paths by  $A(k)$ .

- 3.1.1. Fit multiple regression model for each shortest path** For each shortest path  $e \in A(k)$ , fit the multiple regression model,


$$\mu = 1\beta_0 + X_k \beta_k + X_{e_-} \beta_{e_-} + X_{t(e)} \beta_{t(e)},$$

where  $t(e)$  represents the other terminal vertex of the path  $e$  (other than  $k$ ), and  $e_-$  denotes the index set where the terminal vertex  $t(e)$  is removed from the index set  $e$ . Obtain the  $P$ -value,  $P_{t(e)|\{k, e_-\}}$ , for testing the null hypothesis,  $\beta_{t(e)} = 0$ .

- 3.1.2. Conservative multiple test correction** Declare SNP  $t(e)$  as significant if  $P_{t(e)|\{k, e_-\}} < \alpha / (\alpha_1 p |A(k)|)$  where  $|A(k)|$  is the number of elements of  $A(k)$ .

In Step 3.1.2, the factor  $\alpha_1 p |A(k)|$  can be interpreted as the number of tests accounting for marginal association screening in Step 2. Under the null hypothesis of no effect of all SNPs, the family-wise error rate in detecting any SNPs as declared significant in 3.1.2 is never greater than the nominal level  $\alpha$ : a proof is given in supplementary information Section 3. We propose to declare adjacency between two SNPs  $l$  and  $m$  with correlation  $\rho_{lm}$  if  $a_{\min} \leq \rho_{lm}^2 \leq a_{\max}$  holds. The cutoff  $a_{\min}$  eliminates uncorrelated pairs, and  $a_{\max}$  avoids multicollinearity in the multiple regression fit as described in the Appendix. If there are covariates, we slightly modify the proposed algorithm by adding the covariates to all regression models as with usual covariate adjustment. We investigate appropriate values of  $a_{\min}$  and  $\alpha_1$  in simulation studies where we set  $a_{\max} = 0.8$  throughout. We implemented the proposed method in R (R Core Team, 2015) with input of the result from PLINK's marginal association testing, such as using the `--assoc` option, where the shortest paths in the bidirected graph were collected using the `get.all.shortest.paths` function implemented in the R package `igraph` (Csardi & Nepusz, 2006).





**FIGURE 5** Power with  $m = 1.5$  for six scenarios in quantitative trait simulations with 16 combinations of  $(\alpha_{\min}, \alpha_1) \in \{0.1, 0.3, 0.5, 0.7\} \times \{0.01, 0.001, 0.0001, 0.00001\}$ . Nominal family-wide error rate was set at  $\alpha = 0.05$ . The ordinate is the proportion of runs where any SNP within a 110 kb window upstream of the first causal SNP and downstream of the last causal SNP was significant

number of shortest paths, are given in supplementary information Section 5. These summary measures tended to decrease as  $\alpha_1$  decreased or  $\alpha_{\min}$  increased, and  $\alpha_{\min} = 0.3$  tended to produce longer path lengths than  $\alpha_{\min} = 0.1$ .

### 3.2 | Power

For power simulations, we extracted a region on chromosome 10 ranging from 2,001,233 to 3,499,300 (in NCBI Build 36 base pair position) consisting of 1,273 SNPs. Phenotype data for each sampled subject with these 1,273 SNPs were generated according to a regression model assuming the causal SNPs were the five SNPs listed in Table 1, which also shows the correlation structure. We considered six scenarios with different true regression coefficients based on the values of the  $\beta_j^*$  given in Table 1. We multiplied  $\beta_j^*$  by a constant  $m > 0$  to define the  $j$ th true regression coefficient. Nominal family-wide error rate  $\alpha$  was fixed at 0.05 and total number of SNPs was assumed to be 1,000,000 in all simulations.


#### 3.2.1 | Quantitative traits

We generated by HAPGEN 1,000 simulated datasets of the 1,273 SNPs for  $n = 1000$  subjects, and then we generated phenotype data  $y_i$  for subject  $i$  according to the linear regression model with the five causal SNPs given in Table 1:

$$y_i = \sum_{j=1}^5 X_{i,j} \beta_{0,j} + \epsilon_i,$$

where  $X_{i,j}$  denotes the minor allele count (0, 1, or 2) at the  $j$ th causal SNP ( $j = 1, \dots, 5$ ),  $\beta_{0,j}$  represents the  $j$ th regression coefficient, and  $\epsilon_i$  is an independently and identically distributed standard normal error for the  $i$ th subject. Using each of the six sets of regression coefficients  $\beta_j^*$  in Table 1, we set  $\beta_{0,j} = m\beta_j^*$  for a given constant  $m > 0$ . Two scenarios,  $m = 1$  and  $m = 1.5$ , were considered. Consequently, we carried out 12 power simulations. The proposed method was examined at 16 combinations of the parameters  $(\alpha_{\min}, \alpha_1) \in \{0.1, 0.3, 0.5, 0.7\} \times$





**FIGURE 7** Power with  $m = 1.5$  for six scenarios in binary trait simulations with 16 combinations of  $(a_{\min}, \alpha_1) \in \{0.1, 0.3, 0.5, 0.7\} \times \{0.01, 0.001, 0.0001, 0.00001\}$ . Nominal family-wide error rate was set at  $\alpha = 0.05$ . The ordinate is the proportion of runs where any SNP within a 110 kb window upstream of the first causal SNP and downstream of the last causal SNP was significant

from the general population was generated according to the logistic regression model

$$\mu_i = P(y_i = 1|x_i) = 1 / \left\{ 1 + \exp(-\beta_0 - \sum_{j=1}^5 X_{i,j} \beta_{0,j}) \right\},$$

where  $X_{i,j}$  denotes the minor allele count (0, 1, or 2) at the  $j$ th causal SNP ( $j = 1, \dots, 5$ );  $\beta_{0,j}$  represents the  $j$ th regression coefficient; and  $\beta_0$  is the intercept, which was determined to produce a prevalence of roughly 1%. Using each of the six regression coefficients  $\beta_j^*$  in Table 1, we set  $\beta_{0,j} = 1.4 \times m \beta_j^*$  for a given constant  $m > 0$ . We considered two scenarios,  $m = 1$  and  $m = 1.5$ . We then carried out twelve power simulations. For each replicate, we generated 1,000 subjects with genotype data at these 1,273 SNPs. Subsequently, a binary phenotype was assigned to each subject according to the above logistic regression model. The sampling of 1,000 subjects was repeated until 1,000 cases and 1,000 controls were collected. We simulated 1,000 such datasets. The proposed method was examined at 16

combinations of the parameters  $(a_{\min}, \alpha_1) \in \{0.1, 0.3, 0.5, 0.7\} \times \{0.01, 0.001, 0.0001, 0.00001\}$ . The power of overall detection in the region for  $m = 1$  and  $m = 1.5$  is given in Figures 6 and 7, respectively. The results were roughly consistent with the quantitative power simulations, and the optimal choice of  $(a_{\min}, \alpha_1)$  depended on the underlying genetic architecture.

#### 4 | APPLICATION TO ADNI-GWAS DATA

We applied our proposed method to the ADNI-GWAS dataset obtained from the publicly available data of the Alzheimer’s Disease Neuroimage Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging

**TABLE 2** Significant shortest paths at  $\alpha_{\min} = 0.1$  and  $\alpha_1 = 0.01$ 

Significant path index	CHR	SNP	BP	$P_{\text{marg}}$	$P_{\text{mult}}$
1	19	rs440277	50,053,064	3.8e-03	4.5e-01
1	19	rs387976	50,070,900	8.8e-02	4.1e-02
1	19	rs157580	50,087,106	5.2e-05	2.0e-11
1	19	<b>rs8106922</b>	50,093,506	1.3e-04	1.3e-08
2	19	rs440277	50,053,064	3.8e-03	1.7e-01
2	19	rs387976	50,070,900	8.8e-02	4.3e-02
2	19	rs157580	50,087,106	5.2e-05	1.3e-11
2	19	<b>rs405509</b>	50,100,676	7.7e-04	1.6e-08
4	19	rs157580	50,087,106	5.2e-05	9.2e-11
4	19	<b>rs8106922</b>	50,093,506	1.3e-04	3.9e-10
5	19	rs157580	50,087,106	5.2e-05	1.2e-10
5	19	<b>rs405509</b>	50,100,676	7.7e-04	2.5e-09
8	19	rs8106922	50,093,506	1.3e-04	3.9e-10
8	19	<b>rs157580</b>	50,087,106	5.2e-05	9.2e-11
11	19	rs8106922	50,093,506	1.3e-04	1.2e-01
11	19	rs405509	50,100,676	7.7e-04	1.6e-02
11	19	<b>rs439401</b>	50,106,291	1.3e-05	2.3e-09
12	19	rs405509	50,100,676	7.7e-04	2.5e-09
12	19	<b>rs157580</b>	50,087,106	5.2e-05	1.2e-10
18	19	rs405509	50,100,676	7.7e-04	6.1e-08
18	19	<b>rs439401</b>	50,106,291	1.3e-05	1.2e-09
20	19	rs439401	50,106,291	1.3e-05	5.8e-02
20	19	rs157580	50,087,106	5.2e-05	1.1e-04
20	19	<b>rs8106922</b>	50,093,506	1.3e-04	1.1e-09
21	19	rs439401	50,106,291	1.3e-05	1.2e-09
21	19	<b>rs405509</b>	50,100,676	7.7e-04	6.1e-08

Note. Paths that include genome-wide significant SNPs were excluded. CHR, chromosome; BP, physical position (base-pair);  $P_{\text{marg}}$ , marginal association  $P$ -value;  $P_{\text{mult}}$ , multiple regression  $P$ -value were applied to each path. SNPs in each path are ordered so that the first SNP is adjacent to the phenotype and the last SNP is tested for association. SNPs tested in each path are emphasized in bold.

(MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessments can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). For up-to-date information, see [www.adni-info.org](http://www.adni-info.org). ADNI is an ongoing, longitudinal study with primary purpose being to explore the genetic and neuroimaging information associated with late-onset AD (LOAD). The study investigators recruited elderly subjects older than 65 years of age comprising about 400 subjects with MCI, about 200 subjects with AD, and about 200 healthy controls. Each subject was followed for at least 3 years. During the study period, the subjects were assessed with MRI measures and psychiatric evaluation to determine the diagnosis status at each time point.

The ADNI-GWAS data were obtained from 818 DNA samples of ADNI1 participants using the Illumina Human 610-Quad genotyping array (Shen et al., 2014). The data initially included 620,901 SNPs. We included the *apolipopro-*

*tein E (APOE)* SNPs rs429358 and rs7412 in our analysis. We used data from 684 non-Hispanic Caucasian samples after we excluded one pair showing cryptic relatedness (revealed by the PLINK pairwise  $\hat{\pi}$  statistic being greater than 0.125; Purcell et al., 2007), and we excluded subjects whose reported sex did not match the sex inferred from X-chromosome SNPs. We then applied further quality control measures by excluding SNPs with missing genotype rate  $>0.1$ , Hardy-Weinberg equilibrium test  $P < 10^{-6}$ , and MAF  $< 5\%$ ; the total number of remaining SNPs was 528,984. There were 166 controls and numbers of cases with early MCI (EMCI), late MCI (LMCI), or AD were 67, 110, and 341, respectively. We classified the MCI cases into two subcategories, EMCI and LMCI, following the procedure in the report "Multistate modeling of ADNI progression" that is included with the ADNIMERGE R package provided by ADNI. We then scored the controls and cases of EMCI, LMCI, and AD as 0, 2, 3, and 4, respectively, which was treated

as a quantitative trait in our analysis (we assigned score 1 to cases with significant memory concern [SMC], but the ADNI1 cohort did not contain any subjects with SMC—see <http://adni.loni.usc.edu/study-design/background-rationale/>—and no subjects having score 1 were present in the ADNI-GWAS data). We also considered adjustment for the covariates sex and age in the proposed method. With covariate adjustment,  $P$ -values increased only slightly and the same SNPs were significant, so we only report results without covariate adjustment.

We applied our method with the same 16 combinations of  $(a_{\min}, \alpha_1)$  as in the simulation studies ( $\{0.1, 0.3, 0.5, 0.7\} \times \{0.01, 0.001, 0.0001, 0.00001\}$ ). The nominal family-wise error rate was set as  $\alpha = 0.05$ , in which the number of SNPs, 528,984, was used as the multiple testing correction factor. In Tables 2–5, we show the significant shortest paths and the SNPs contained in them. We do not show shortest paths containing SNPs with genome-wide significance (i.e., those with marginal association  $P$ -value less than  $0.05/528,984$ ), so that only SNPs undetected by marginal association scans are shown (a full list of results is given in supplementary information Section 5). Without paths containing genome-wide significant SNPs, four combinations of  $(a_{\min}, \alpha_1)$  remained:  $(0.1, 0.01)$ ,  $(0.1, 0.001)$ ,  $(0.1, 0.0001)$ , and  $(0.3, 0.001)$ . All SNPs detected with  $(0.1, 0.0001)$  were also contained among the significant results with  $(0.1, 0.01)$  and  $(0.1, 0.001)$ , which implies that  $\alpha_1 = 0.0001$  is too stringent a threshold, as was observed in our simulation studies. The ADNI-GWAS application also implies that  $a_{\min} = 0.7$  is unlikely to provide high power, again as observed in our simulation studies.

As observed in Tables 2–5, SNPs with weak marginal signals showed enhanced association signals with multiple regression. The results varied with the choice of parameters  $(a_{\min}, \alpha_1)$ , but SNPs on chromosome 19 were detected in most of the scenarios. The noteworthy regions contain *APOE4* SNPs, which are also genome-wide significant even with the marginal association test, SnipSnip, and UNPHASED, as shown in Table 6. We also applied lasso (Tibshirani, 1996) using the PUMA software (Hoffman, Logsdon, & Mezey, 2013) with the default setting and optimal model chosen according to Akaike information criterion. The SNPs detected with lasso are shown in supplementary information Section 5 along with  $P$ -values from PUMA (PUMA  $P$ -values are a ranking measure, not the usual measure of statistical significance). Six unique SNPs on chromosome 5 given in Tables 3 and 5 are more interesting, as they were not detected by the marginal association test, SnipSnip, UNPHASED, or lasso. The gene closest to these unique SNPs on chromosome 5 is *semaphorin 5A* (*SEMA5A*). The Ensembl Variant Effect Predictor (McLaren et al., 2010) predicted that these SNPs are intergenic and function as modifiers. The *SEMA5A* gene was previously reported to be a susceptibility locus for

**TABLE 3** Significant shortest paths at  $a_{\min} = 0.1$  and  $\alpha_1 = 0.001$

Significant path index	CHR	SNP	BP	$P_{\text{marg}}$	$P_{\text{mult}}$
1	5	rs2963337	8,988,632	1.5e-04	4.0e-08
1	5	rs2963345	8,962,508	3.2e-03	7.0e-03
1	5	<b>rs6870451</b>	8,991,885	9.3e-02	8.6e-07
2	5	rs2963337	8,988,632	1.5e-04	3.8e-08
2	5	rs2963345	8,962,508	3.2e-03	7.9e-03
2	5	<b>rs11955429</b>	8,994,715	7.8e-02	9.9e-07
4	19	rs157580	50,087,106	5.2e-05	9.2e-11
4	19	<b>rs8106922</b>	50,093,506	1.3e-04	3.9e-10
5	19	rs157580	50,087,106	5.2e-05	1.2e-10
5	19	<b>rs405509</b>	50,100,676	7.7e-04	2.5e-09
8	19	rs8106922	50,093,506	1.3e-04	3.9e-10
8	19	<b>rs157580</b>	50,087,106	5.2e-05	9.2e-11
11	19	rs8106922	50,093,506	1.3e-04	1.2e-01
11	19	rs405509	50,100,676	7.7e-04	1.6e-02
11	19	<b>rs439401</b>	50,106,291	1.3e-05	2.3e-09
12	19	rs405509	50,100,676	7.7e-04	2.5e-09
12	19	<b>rs157580</b>	50,087,106	5.2e-05	1.2e-10
18	19	rs405509	50,100,676	7.7e-04	6.1e-08
18	19	<b>rs439401</b>	50,106,291	1.3e-05	1.2e-09
20	19	rs439401	50,106,291	1.3e-05	5.8e-02
20	19	rs157580	50,087,106	5.2e-05	1.1e-04
20	19	<b>rs8106922</b>	50,093,506	1.3e-04	1.1e-09
21	19	rs439401	50,106,291	1.3e-05	1.2e-09
21	19	<b>rs405509</b>	50,100,676	7.7e-04	6.1e-08

Note. Paths that include genome-wide significant SNPs were excluded. CHR, chromosome; BP, physical position (base-pair);  $P_{\text{marg}}$ , marginal association  $P$ -value;  $P_{\text{mult}}$ , multiple regression  $P$ -value applied to each path. SNPs in each path are ordered so that the first SNP is adjacent to the phenotype and the last SNP is tested for association. SNPs tested in each path are emphasized in bold.

Parkinson's disease (Lin, Lesnick, Maraganore, & Isacson, 2009; Maraganore et al., 2005) and for autism (Melin et al., 2006). *SEMA5A* is involved in axonal guidance during neural development.

## 5 | DISCUSSION

We present herein a new conservative multiple testing method for detecting genetic association signals hidden by LD. Our proposed bidirected graph approach can be applied to high-dimensional genome-wide SNP data. Despite the conservativeness of our proposed method, simulation studies confirmed that the method has higher power than existing methods. In an application to ADNI-GWAS data, our method detected new susceptibility SNPs on chromosome 5 that were originally undetected by existing methods, including lasso. The negative results with lasso are expected because

**TABLE 4** Significant shortest paths at  $a_{\min} = 0.1$  and  $\alpha_1 = 0.0001$ 

Significant path index	CHR	SNP	BP	$P_{\text{marg}}$	$P_{\text{mult}}$
2	19	rs157580	50,087,106	5.2e-05	9.2e-11
2	19	<b>rs8106922</b>	50,093,506	1.3e-04	3.9e-10
3	19	rs157580	50,087,106	5.2e-05	1.2e-10
3	19	<b>rs405509</b>	50,100,676	7.7e-04	2.5e-09
7	19	rs439401	50,106,291	1.3e-05	5.8e-02
7	19	rs157580	50,087,106	5.2e-05	1.1e-04
7	19	<b>rs8106922</b>	50,093,506	1.3e-04	1.1e-09
8	19	rs439401	50,106,291	1.3e-05	1.2e-09
8	19	<b>rs405509</b>	50,100,676	7.7e-04	6.1e-08

Note. Paths that include genome-wide significant SNPs were excluded. CHR, chromosome; BP, physical position (base-pair);  $P_{\text{marg}}$ , marginal association  $P$ -value;  $P_{\text{mult}}$ , multiple regression  $P$ -value applied to each path. SNPs in each path are ordered so that the first SNP is adjacent to the phenotype and the last SNP is tested for association. SNPs tested in each path are emphasized in bold.

**TABLE 5** Significant shortest paths at  $a_{\min} = 0.3$  and  $\alpha_1 = 0.001$ 

Significant path index	CHR	SNP	BP	$P_{\text{marg}}$	$P_{\text{mult}}$
1	5	rs2963327	8,987,159	2.0e-04	8.1e-08
1	5	rs2963345	8,962,508	3.2e-03	7.1e-03
1	5	<b>rs6870451</b>	8,991,885	9.3e-02	1.4e-06
2	5	rs2963327	8,987,159	2.0e-04	7.8e-08
2	5	rs2963344	8,962,428	3.7e-03	1.1e-02
2	5	<b>rs6870451</b>	8,991,885	9.3e-02	2.0e-06
3	5	rs2963327	8,987,159	2.0e-04	7.7e-08
3	5	rs2963345	8,962,508	3.2e-03	8.0e-03
3	5	<b>rs11955429</b>	8,994,715	7.8e-02	1.6e-06
4	5	rs2963327	8,987,159	2.0e-04	7.0e-08
4	5	rs2963344	8,962,428	3.7e-03	1.3e-02
4	5	<b>rs11955429</b>	8,994,715	7.8e-02	2.3e-06
5	5	rs2963337	8,988,632	1.5e-04	4.0e-08
5	5	rs2963345	8,962,508	3.2e-03	7.0e-03
5	5	<b>rs6870451</b>	8,991,885	9.3e-02	8.6e-07
6	5	rs2963337	8,988,632	1.5e-04	3.9e-08
6	5	rs2963344	8,962,428	3.7e-03	1.1e-02
6	5	<b>rs6870451</b>	8,991,885	9.3e-02	1.3e-06
7	5	rs2963337	8,988,632	1.5e-04	3.8e-08
7	5	rs2963345	8,962,508	3.2e-03	7.9e-03
7	5	<b>rs11955429</b>	8,994,715	7.8e-02	9.9e-07
8	5	rs2963337	8,988,632	1.5e-04	3.5e-08
8	5	rs2963344	8,962,428	3.7e-03	1.2e-02
8	5	<b>rs11955429</b>	8,994,715	7.8e-02	1.4e-06

Note. Paths that include genome-wide significant SNPs were excluded. CHR, chromosome; BP, physical position (base-pair);  $P_{\text{marg}}$ , marginal association  $P$ -value;  $P_{\text{mult}}$ , multiple regression  $P$ -value applied to each path. SNPs in each path are ordered so that the first SNP is adjacent to the phenotype and the last SNP is tested for association. SNPs tested in each path are emphasized in bold.

lasso is a variant of stagewise regression (Efron, Hastie, Johnstone, & Tibshirani, 2004); in view of lasso's iterative model update process (c.f. the coordinate descent algorithm; Friedman, Hastie, & Tibshirani, 2010), an SNP is entered into the model only if the absolute value of the correlation between

that SNP and the residual of the current model is larger than some threshold as achieved by the soft-thresholding function. This in turn implies that an SNP weakly correlated with the current residual will never be included in the model. Hidden SNPs having large effects conditional on other SNPs will

**TABLE 6** SNPs that were significant by the marginal association test, SnipSnip, and UNPHASED; family-wise error rate was 5% using Bonferroni correction based on the number of SNPs, 528,984

CHR	SNP	BP	$P_{\text{marg}}$	$P_{\text{SnipSnip}}$	$P_{\text{Unphased}}$
19	rs387976	50,070,900	8.9e-02	6.0e-01	1.5e-09
19	rs11667640	50,071,631	2.0e-02	3.4e-02	1.9e-12
19	rs6859	50,073,874	1.9e-02	9.4e-02	5.8e-11
19	rs157580	50,087,106	5.2e-05	1.2e-10	1.0e-18
19	rs2075650	50,087,459	1.9e-11	6.3e-09	5.3e-19
19	rs8106922	50,093,506	1.3e-04	3.9e-10	5.0e-21
19	rs405509	50,100,676	7.7e-04	2.5e-09	2.5e-22
19	rs429358	50,103,781	9.5e-23	1.5e-18	5.6e-20

CHR, chromosome; BP, physical position (base-pair).

be included only if the current model contains those other SNPs. However, if the conditioning SNPs have only weak or moderately sized marginal signals, noninformative SNPs will be entered by chance into the model long before the conditioning SNPs are entered. Consequently, the noninformative SNPs result in a model with unnecessarily high dimension, wasting degrees of freedom, and model building will terminate before the important, hidden SNP is included. In contrast, our method avoids including too many noninformative SNPs among the conditioning SNPs by restricting to the shortest paths in a “local” bidirected graph, making the hypothesis test more powerful.

Our theoretical consideration of the presence of multiple causal variants that are in LD is also new. So far, designing GWASs (in particular, selecting tag SNPs) has been based chiefly on statistical power analyses that assume causal variants exist separately in unlinked loci (de Bakker et al., 2005; Eberle et al., 2007; Spencer et al., 2009). Conclusions from existing studies based on the GWAS design may therefore not be generalizable to situations with multiple causal variants that are in LD. One consequence is that current SNP panels are suboptimal in this situation; hence, additional studies of statistical power are needed. However, power calculations turn out to be much more complicated than in situations with multiple causal variants that are not in LD: it remains unknown which of various genetic models and statistical methods is optimal.

Although our proposed method attempts to detect hidden SNPs that are difficult to detect with GWAS data (Eberle et al., 2007), detection should be distinguished from true association. Even though a particular combination of detected SNPs produces the highest power, it does not necessarily indicate that these are the causal SNPs. Subsequent, careful examination based on biological background and function is also necessary.

There are several possible directions for future work to improve our method. First, our proposed test is slightly conservative as shown in type I error simulations. This is because the method is based on the Bonferroni-type control of multiplicity

of shortest paths. There is inevitably overlap in the SNPs to be tested, resulting in redundancy. Development of a less conservative method of type I error control may further improve the statistical power. Second, our method has two tuning parameters ( $a_{\text{min}}, \alpha_1$ ); although the choice influences the result, we could not determine an optimal choice that could be used in all simulation scenarios. Because the optimal parameters seemed to depend on underlying genetic architecture, which is unknown, we recommend trying multiple sets of ( $a_{\text{min}}, \alpha_1$ ) in practice. Simulation studies suggested that the two parameters be in the ranges  $a_{\text{min}} \in [0.1, 0.5]$  and  $\alpha_1 \in [0.01, 0.001]$ , which produced improved performance over existing methods. Application to the ADNI-GWAS data also supported this choice of ranges. Based on our experience with applications to other real GWAS data (not shown), we found that  $a_{\text{min}} = 0.3$  and  $\alpha_1 = 0.001$  tended to allow detection of SNPs that had gone undetected with existing methods.

Our method builds a bidirected graph locally within a fixed-size window centered (focused) around each SNP across the genome. This approach may fail in the presence of long-distance LD, which is not ignorable in real human data (Pritchard & Przeworski, 2001). However, use of a wider window size incurs both computational and multiple testing burdens as the number of shortest paths increases exponentially. As observed in simulation studies, the LD between two SNPs needs to be high for them to be adjacent in the bidirected graph. The cutoff  $a_{\text{min}}$  approximately corresponds to that for  $r^2$ . According to the relationship between average  $r^2$  and genetic distance in human data given in Figure 2 of (Pritchard & Przeworski, 2001), the average of  $r^2$  is much lower than 0.1 for a 0.1 cM distance between SNPs. Thus, our choice of a 100 kb window could be reasonable on average in practice, although exceptions surely exist. We will continue applying our method to real human genetic data to explore the optimal choices of the parameters in future research. R code that implements our method for SNP-GWAS data in PLINK binary format is available from the authors upon request.

## ACKNOWLEDGMENTS

The authors thank a referee, Prof. Shete, and Dr. Cologne for helpful comments that led to significant improvement of the paper. Data collection and sharing for this project was funded by the Alzheimer’s Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer’s Association; Alzheimer’s Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Eisai, Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company;

EuroImmun; F. Hoffmann-La Roche Ltd. and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC; Johnson & Johnson Pharmaceutical Research & Development LLC; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer, Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health ([www.fnih.org](http://www.fnih.org)). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California. This work was carried out under the ISM General Cooperative Research 1 (2015-ISM-CRP-1013). This work was supported by JSPS KAKENHI grant numbers JP16K00064, JP16K08638, JP16H05242, JP15H01678, JP16H01528.

## REFERENCES

- Cox, D. R., & Wermuth, N. (1996). *Multivariate dependencies*. London: Chapman and Hall.
- Csardi, G., & Nepusz, T. (2006). The igraph software package for complex network research. *InterJournal, Complex Systems*, 1695. <http://igraph.org>
- de Bakker, P. I., Yelensky, R., Pe'er, I., Gabriel, S. B., Daly, M. J., & Altshuler, D. (2005). Efficiency and power in genetic association studies. *Nature Genetics*, 37, 1217–1223.
- Drton, M., & Perlman, M. (2007). Multiple testing and error control in Gaussian graphical model selection. *Statistical Science*, 22, 430–449.
- Dudbridge, F. (2008). Likelihood-based association analysis for nuclear families and unrelated subjects with missing genotype data. *Human Heredity*, 66, 87–98.
- Dudbridge, F., Holmans, P. A., & Wilson, S. G. (2011). A flexible model for association analysis in sibships with missing genotype data. *Annals of Human Genetics*, 75, 428–438.
- Eberle, M. A., Ng, P. C., Kuhn, K., Zhou, L., Peiffer, D. A., Galver, L., ... Murray, S. S. (2007). Power to detect risk alleles using genome-wide tag SNP panels. *PLoS Genetics*, 3, e170.
- Efron, B., Hastie, T., Johnstone, I., & Tibshirani, R. (2004). Least angle regression. *Annals of Statistics*, 32, 407–499.
- Ehret, G. B., Lamparter, D., Hoggart, C. J., Genetic Investigation of Anthropometric Traits Consortium, Whittaker, J. C., Beckmann, J. S., Kutalik, Z. (2012). A multi-SNP locus-association method reveals a substantial fraction of the missing heritability. *American Journal of Human Genetics*, 91, 863–871.
- Friedman, J., Hastie, T., & Tibshirani, R. (2010). Regularization paths for generalized linear models via coordinate descent. *Journal of Statistical Software*, 33, 1–22.
- Hoffman, G. E., Logsdon, B. A., & Mezey, J. G. (2013). PUMA: A unified framework for penalized multiple regression analysis of GWAS data. *PLoS Computational Biology*, 9, e1003101.
- Howey, R., & Cordell, H. J. (2014). Imputation without doing imputation: A new method for the detection of non-genotyped causal variants. *Genetic Epidemiology*, 38, 173–190.
- Kauermann, G. (1996). On a dualization of graphical Gaussian models. *Scandinavian Journal of Statistics*, 23, 105–116.
- Kim, S., Morris, N., Won, S., & Elston, R. C. (2010). Single-marker and two-marker association tests for unphased case-control genotype data, with a power comparison. *Genetic Epidemiology*, 34, 66–77.
- Lin, L., Lesnick, T. G., Maraganore, D. M., & Isacson, O. (2009). Axon guidance and synaptic maintenance: Preclinical markers for neurodegenerative disease and therapeutics. *Trends in Neurosciences*, 32, 142–149.
- Maher, B. (2008). Personal genomes: The case of the missing heritability. *Nature*, 456, 18–21.
- Manolio, T. A. (2013). Bringing genome-wide association findings into clinical use. *Nature Reviews Genetics*, 14, 549–558.
- Manolio, T. A., Collins, F. S., Cox, N. J., Goldstein, D. B., Hindorf, L. A., Hunter, D. J., ... Visscher, P. M. (2009). Finding the missing heritability of complex diseases. *Nature*, 461, 747–753.
- Maraganore, D. M., de Andrade, M., Lesnick, T. G., Strain, K. J., Farrer, M. J., Rocca, W. A., ... Ballinger, D. G. (2005). High-resolution whole-genome association study of Parkinson disease. *American Journal of Human Genetics*, 77, 685–693.
- McLaren, W., Pritchard, B., Rios, D., Chen, Y., Flicek, P., & Cunningham, F. (2010). Deriving the consequences of genomic variants with the Ensembl API and SNP Effect Predictor. *Bioinformatics*, 26, 2069–2070.
- Melin, M., Carlsson, B., Anckarsater, H., Rastam, M., Betancur, C., Isaksson, A., ... Dahl, N. (2006). Constitutional downregulation of SEMA5A expression in autism. *Trends in Neurosciences*, 54, 64–69.
- Pritchard, J. K., & Przeworski, M. (2001). Linkage disequilibrium in humans: Models and data. *American Journal of Human Genetics*, 69, 1–14.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., ... Sham, P. C. (2007). PLINK: A tool set for whole-genome association and population-based linkage analyses. *American Journal of Human Genetics*, 81, 559–575.
- R Core Team. (2015). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rao, C. R., & Yanai, H. (1979). General definition and decomposition of projectors and some applications to statistical problems. *Journal of Statistical Planning and Inference*, 3, 1–17.
- Shen, L., Thompson, P. M., Potkin, S. G., Bertram, L., Farrer, L. A., Foroud, T. M., ... Alzheimer's Disease Neuroimaging Initiative. (2014). Genetic analysis of quantitative phenotypes in AD and MCI: Imaging, cognition and biomarkers. *Brain Imaging and Behavior*, 8, 183–207.
- Slavin, T. P., Feng, T., Schnell, A., Zhu, X., & Elston, R. C. (2011). Two-marker association tests yield new disease associations for coronary artery disease and hypertension. *Human Genetics*, 130, 725–733.

- Spencer, C. C. A., Su, Z., Donnelly, P., & Marchini, J. (2009). Designing genome-wide association studies: Sample size, power, imputation, and the choice of genotyping chip. *PLoS Genetics*, *5*, e1000477.
- Su, Z., Marchini, J., & Donnelly, P. (2011). HAPGEN: Simulation of multiple disease SNPs. *Bioinformatics*, *5*, 2304–2305.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society Series B*, *58*, 267–288.
- Ueki, M., & Kawasaki, Y. (2013). Multiple choice from competing regression models under multicollinearity based on standardized update. *Computational Statistics & Data Analysis*, *63*, 31–41.
- Wang, X., Morris, N. J., Schaid, D. J., & Elston, R. C. (2012). Power of single- vs. multi-marker tests of association. *Genetic Epidemiology*, *36*, 480–487.

- Wellek, S., & Ziegler, A. (2009). A genotype-based approach to assessing the association between single nucleotide polymorphisms. *Human Heredity*, *67*, 128–139.

## SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

**How to cite this article:** Ueki M, Kawasaki Y, Tamiya G. Detecting genetic association through shortest paths in a bidirected graph. *Genet. Epidemiol.* 2017;41:481–497. <https://doi.org/10.1002/gepi.22051>

## APPENDIX

### Motivation for the bidirected graph approach

Here, we describe the theoretical background behind our proposed method. First, we relate the bidirected graph with the least-squares estimator  $\hat{\beta}_{j,A}$  in the model  $y = x_j\beta_j + \sum_{k \in A} x_k\beta_k + \epsilon$ . Let  $X_A = (x_k : k \in A)$ . To this end, we exploit the following representation (Rao & Yanai, 1979; Ueki & Kawasaki, 2013):

$$\hat{\beta}_{j,A} = \|\mathcal{Q}_{X_A} x_j\|^{-2} y^T (\mathcal{Q}_{X_A} x_j).$$

Its expectation and variance are given by

$$E(\hat{\beta}_{j,A}|X) = \|\mathcal{Q}_{X_A} x_j\|^{-2} \mu^T (\mathcal{Q}_{X_A} x_j)$$

and  $\text{var}(\hat{\beta}_{j,A}|X) = \sigma^2 \|\mathcal{Q}_{X_A} x_j\|^{-2}$ , respectively. Consequently, in order to evaluate whether  $E(\hat{\beta}_{j,A}|X) = 0$  or not, it suffices to investigate the quantity

$$\mu^T (\mathcal{Q}_{X_A} x_j).$$

The following result, whose proof is given in supplementary information Section 1, helps us to screen SNPs to be entered into the conditioning set  $A$  for  $\hat{\beta}_{j,A}$ . It is essentially the equivalence between pairwise and global Markov properties in a bidirected graph for a multivariate Gaussian distribution (Drton & Perlman, 2007; Kauermann, 1996).

**PROPOSITION 1.** *If there is no path between vertexes 0 (i.e.,  $\mu = E(y|X)$ ) and  $j \in M = V \setminus \{0\}$  with intermediate vertexes being any subset from a vertex set  $\{j_1, \dots, j_l\} \subset M \setminus \{j\} = V \setminus \{0, j\}$  (allowing the empty set), then for any  $A \subset \{j_1, \dots, j_l\}$ , we have  $\mu^T (\mathcal{Q}_{X_A} x_j) = 0$ .*

Proposition 1 implies that, for the discovery of SNP  $j$  such that  $E(\hat{\beta}_{j,A}|X) \neq 0$  for some  $A$ , we can ignore the SNPs having no paths to  $\mu = E(y|X)$ . Conversely, we can focus on SNPs having paths to SNPs that are adjacent to  $\mu$ . The adjacency with  $\mu$  corresponds to the presence of a marginal association signal as in a standard GWAS scan. In typical GWAS, however, SNPs often show weak marginal signals. Our target is the SNPs with low marginal association signals. To this end, we focus on SNPs that show a moderate marginal signal (e.g. marginal  $P$ -value less than a cutoff), and then build a bidirected graph around each of those SNPs. In samples from the general population, LD between two SNPs far apart is expected to be low. We exploit this LD structure in application to GWASs. Namely, it suffices to consider SNPs for the conditioning set  $A$  that are close to the focal SNP to be tested for the effect; SNPs faraway can be excluded from the candidates for  $A$ . As a consequence, we build a bidirected graph locally, for example, 100 kb (kilobase pair) window both upstream and downstream of each SNP.

We note that Proposition 1 does not guarantee its complementary statement—that is, the existence of a path between  $\mu$  (vertex 0) and SNP  $j$  with intermediate vertexes  $\{j_1, \dots, j_l\}$  implies the existence of some subset  $A \subset \{j_1, \dots, j_l\}$  such that  $E(\hat{\beta}_{j,A}|X) \neq 0$ . In fact, the complement of Proposition 1 does not always hold. For instance, if there are two distinct vertexes  $j$

and  $k$  whose corresponding SNPs satisfy  $x_j = x_k = \mu$ , then there is a path between vertexes 0 and  $j$  with an intermediate vertex  $k$ . We can easily obtain the result that  $\mu^T(Q_{x_k} x_j) = 0$  as  $Q_{x_k} x_j$  is the residual of  $x_j$  from regression onto  $x_k$ . More generally, if  $x_j$  is represented by a linear combination of  $X_A$  (i.e., there is a vector  $\gamma_A \in \mathbb{R}^{|A|}$  such that  $x_j = X_A \gamma_A$ ), then  $\mu^T Q_{X_A} x_j = 0$ . From the relationship with the regression coefficient of  $x_j$  in multiple regression of  $\mu$  on  $(x_j, X_A)$ , this corresponds to perfect multicollinearity. Consequently, we still have the possibility that  $\mu^T Q_{X_A} x_j = 0$  using the SNPs on the paths to  $\mu$  with terminal vertex  $j$  and intermediate vertexes  $A$ . In what follows, we explore a way of eliminating such redundant paths.

In the causal-inference literature, assumptions such as faithfulness are usually imposed to ensure that the complement of Proposition 1 is true. However, rather than assuming such conditions, we explore conditions under which the complementary result of Proposition 1 holds. Define the length of the path between vertexes  $a$  and  $b$  with intermediate vertexes  $\{j_1, \dots, j_l\}$  by the number of edges on the path, i.e.,  $l$ . The length between two adjacent vertexes  $a$  and  $b$  is defined to be 0. We declare that the path between vertexes  $a$  and  $b$  with length  $l \geq 0$  is the shortest if and only if there is no path between  $a$  and  $b$  whose length is less than  $l$ . This definition permits the existence of multiple shortest paths with equal length.

We then have the following result, which complements Proposition 1.

**PROPOSITION 2.** *Suppose that there is at least one path between vertexes 0 and  $j$ , and that the set of intermediate vertexes of one of the shortest paths is  $\{j_1, \dots, j_l\} \subset M \setminus \{j\} = V \setminus \{0, j\}$ . Then, we have  $\mu^T(Q_{X_{\{j_1, \dots, j_l\}}} x_j) \neq 0$ .*

The proof is given in supplementary information Section 2. For the discovery of SNP  $j$  such that  $E(\hat{\beta}_{j,A}|X) \neq 0$  for some  $A$ , Proposition 2 suggests that we should incorporate into the conditioning set  $A$  the intermediate vertexes of the shortest paths between  $\mu = E(y|X)$  and  $j$ . In the above example for two distinct vertexes  $j$  and  $k$  whose corresponding SNPs satisfy  $x_j = x_k = \mu$ , the length of the path between vertexes 0 and  $j$  with an intermediate vertex  $k$  is one, which is not the shortest because vertexes 0 and  $j$  are adjacent. That is, the path between 0 and  $j$  (whose length is 0) is the shortest path. Proposition 2 states that  $\mu^T x_j \neq 0$  (the shortest path) is guaranteed but  $\mu^T Q_{x_k} x_j \neq 0$  is not guaranteed (not the shortest path). Note that Proposition 2 does not eliminate the possibility that the nonshortest path gives  $\mu^T Q_A x_j \neq 0$ . Therefore, it is possible that our proposed method misses some nonshortest paths that give a large nonzero effect. Nevertheless, we consider that restricting the search to the shortest paths is useful because the shortest paths produce parsimonious regression models with low degrees of freedom. This narrows the candidate models to be considered, thereby improving statistical power.

Because Propositions 1 and 2 are based on the bidirected graph with which adjacency is based on the Wellek-Ziegler correlation (which can take value 0), it is not practical in real data applications. To resolve this, we introduce cutoff values for the Wellek-Ziegler correlation between SNPs to declare adjacency in the bidirected graph. Specifically, for given correlation cutoffs  $a_{\min}$  and  $a_{\max}$ , if  $(x_j^T x_k)^2 \in [a_{\min}, a_{\max}] \subset [0, 1]$ , we declare that SNPs  $j$  and  $k$  are adjacent; otherwise, we declare that they are not adjacent. We now derive  $a_{\max}$  empirically as follows. Consider a simple multicollinear case of two SNPs  $x_1$  and  $x_2$  such that  $x_1^T x_2 \approx 1$  and  $x_2^T \mu \neq 0$ , and hence the residual  $Q_{x_2} x_1 \approx 0$ . Then, on the path  $\mu \leftrightarrow x_2 \leftrightarrow x_1$ , we will have that  $x_1^T Q_{x_2} \mu \approx 0$ , that is, the effect of  $x_1$  has been removed by  $x_2$ . Because the test statistic for testing the null hypothesis  $\beta_1 = 0$  in the regression model  $\mu = x_1 \beta_1 + x_2 \beta_2$  is proportional to the correlation between  $y$  and  $Q_{x_2} x_1$  (Remark 1 in Ueki & Kawasaki, 2013), the inclusion of  $x_2$ , which is highly correlated with  $x_1$ , renders the test underpowered due to the above-mentioned effect removal. A similar consideration can be found in Howey and Cordell (2014) who noted that the power of their proposed artificial imputation test tends to improve by choosing a partner SNP such that the correlation with the anchor SNP is neither high nor low. Even with multiple  $x_2$ , the test statistic for testing the null hypothesis  $\beta_1 = 0$  is proportional to the correlation between  $y$  and  $Q_{x_2} x_1$  (Ueki & Kawasaki, 2013), so multicollinearity leads to underpowered detection due to effect removal. In that case, the cutoff  $a_{\max}$  avoids the inclusion of highly correlated SNPs simultaneously, and it is expected that multicollinearity will be alleviated.