

## CSF amyloid is a consistent predictor of white matter hyperintensities across the disease course from aging to Alzheimer's disease



Phoebe Walsh<sup>a,\*</sup>, Carole H. Sudre<sup>a,b,c</sup>, Cassidy M. Fiford<sup>a</sup>, Natalie S. Ryan<sup>a</sup>, Tammarny Lashley<sup>d,e</sup>, Chris Frost<sup>a,f</sup>, Josephine Barnes<sup>a</sup>, For the ADNI Investigators<sup>1</sup>

<sup>a</sup> Dementia Research Centre, Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, London, UK

<sup>b</sup> Department of Biomedical Engineering, School of Biomedical Engineering and Imaging Sciences, King's College London, London, UK

<sup>c</sup> Centre for Medical Image Computing, University College London, London, UK

<sup>d</sup> Queen Square Brain Bank for Neurological Disorders, Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, London, UK

<sup>e</sup> Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, London, UK

<sup>f</sup> Department of Medical Statistics, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK

### ARTICLE INFO

#### Article history:

Received 22 July 2019

Received in revised form 5 March 2020

Accepted 10 March 2020

Available online 17 March 2020

#### Keywords:

Alzheimer's disease

Vascular disease

White matter hyperintensities

Amyloid

Tau

Cerebrospinal fluid

### ABSTRACT

This study investigated the relationship between white matter hyperintensities (WMH) and cerebrospinal fluid (CSF) Alzheimer's disease (AD) biomarkers. Subjects included 180 controls, 107 individuals with a significant memory concern, 320 individuals with early mild cognitive impairment, 171 individuals with late mild cognitive impairment, and 151 individuals with AD, with 3T MRI and CSF A $\beta$ 1–42, total tau (t-tau), and phosphorylated tau (p-tau) data. Multiple linear regression models assessed the relationship between WMH and CSF A $\beta$ 1–42, t-tau, and p-tau. Directionally, a higher WMH burden was associated with lower CSF A $\beta$ 1–42 within each diagnostic group, with no evidence for a difference in the slope of the association across diagnostic groups ( $p = 0.4$ ). Pooling all participants, this association was statistically significant after adjustment for t-tau, p-tau, age, diagnostic group, and APOE- $\epsilon$ 4 status ( $p < 0.001$ ). Age was the strongest predictor of WMH (partial  $R^2 \sim 16\%$ ) compared with CSF A $\beta$ 1–42 (partial  $R^2 \sim 5\%$ ). There was no evidence for an association with WMH and either t-tau or p-tau. These data are supportive of a link between amyloid burden and presumed vascular pathology.

© 2020 Elsevier Inc. All rights reserved.

### 1. Introduction

The co-existence of cerebrovascular disease (CVD) and Alzheimer's disease (AD) is increasingly well recognized; patients with AD often display both AD pathology and underlying vascular pathologies at autopsy such as small vessel disease (SVD), microvascular injury, and cerebrovascular lesions (Brayne et al., 2009; Esiri et al., 2014; Jellinger and Attems, 2006; Kapasi et al., 2017;

Schneider et al., 2007, 2009). Cerebrovascular pathology is well established as an important contributor to cognitive decline, acting as the primary cause of at least 20% of dementias (Gorelick et al., 2011). However, the relationship between CVD and the hallmark Alzheimer's pathologies including tau and amyloid beta (A $\beta$ ) is yet to be fully characterized.

A $\beta$  deposition in the brain can be globally measured in vivo by analyzing cerebrospinal fluid (CSF) levels of A $\beta$ 1–42, whereby an increase in parenchymal A $\beta$  is signified by a decrease in CSF A $\beta$ 1–42 (Blennow et al., 2010). Cerebral tau levels can also be assessed using CSF analysis, with the levels of total tau (t-tau) and phosphorylated tau (p-tau) thought to indicate neuronal damage and pathological tau deposition, respectively (Jack et al., 2018). Markers of CVD are visible and quantifiable on magnetic resonance imaging (MRI), with perhaps the best characterized being white matter hyperintensities (WMH) that appear hyperintense on T2-weighted or FLAIR imaging and hypointense on T1-weighted imaging (Wardlaw et al., 2013). WMH have numerous histopathological correlates, including

\* Corresponding author at: Dementia Research Centre, UCL Queen Square Institute of Neurology, Russell Square House, 10–12 Russell Square, London WC1B 5EH, UK. Tel.: 0203 1086 222; fax: 0203 448 3104

E-mail address: [phoebe.walsh.10@ucl.ac.uk](mailto:phoebe.walsh.10@ucl.ac.uk) (P. Walsh).

<sup>1</sup> Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database ([adni.loni.usc.edu](http://adni.loni.usc.edu)). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: [http://adni.loni.usc.edu/wp-content/uploads/how\\_to\\_apply/ADNI\\_Acknowledgement\\_List.pdf](http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf).

ependymal loss, cerebral ischemia, and demyelination (Gouw et al., 2011). However, it may be that WMH are not only a marker of CVD, but also occur as a result of Alzheimer's pathologies (McAleese et al., 2015, 2017).

Research studies are now moving toward a more biomarker led approach to AD (Jack et al., 2018), making it important to understand how possible biomarkers are interrelated at different stages of disease. Although no cerebrovascular biomarkers have been included in the research framework proposed by Jack et al. (2018), WMH are a good candidate marker to enhance our understanding of the associations between AD and vascular pathology.

WMH have shown associations with amyloid accumulation, as measured by positron emission tomography (Kandel et al., 2016; Marnane et al., 2016) and CSF biomarkers (Marnane et al., 2016; Pietrobboni et al., 2018). Tau aggregation has also been linked to WMH burden (McAleese et al., 2015, Tosto et al., 2015), although results are somewhat less consistent (Kester et al., 2014; Osborn et al., 2018). The aim of this study was to add evidence to further characterize these complex relationships between AD and vascular biomarkers, and to extend the current literature by examining these relationships across the full spectrum from normal aging to AD. We used continuous as opposed to binary measures of AD biomarkers, to more sensitively examine relationships with WMH. To the best of our knowledge, this study is the first to look at relationships between traditional biomarkers of AD (CSF A $\beta$ 1-42, t-tau, and p-tau) and WMH in controls, individuals with a significant memory concern (SMC), individuals with early MCI (EMCI), individuals with late MCI (LMCI), and individuals with AD. We hypothesize that WMH will show associations with CSF A $\beta$ 1-42, t-tau, and p-tau biomarkers, reflecting the potential relationships between vascular dysfunction, amyloid deposition, neurodegeneration, and tau deposition. Because WMH are likely to represent different admixtures of pathology at different disease stages, with recent evidence suggesting in later disease stages WMH associations with tau may be stronger (McAleese et al., 2017) and A $\beta$ 1-42 may be weaker (Zhou et al., 2009), we also hypothesize that the relationships between A $\beta$  and tau biomarkers and WMH may change throughout the disease course with stronger associations between tau markers and WMH at later disease stages, and weaker associations between A $\beta$ 1-42 and WMH at the more advanced stages.

## 2. Materials and methods

### 2.1. Cohort

The demographic, imaging, and biological data used for this study were downloaded from the ADNI database (<http://adni.loni.usc.edu/>). Both ADNIGO and ADNI2 data were used. ADNI is a multicenter, longitudinal public-private funded partnership, with the primary goal of using demographic, biomarker, neuropsychological, and MRI data to monitor progression of AD. Since 2003, Principle Investigator Michael W. Weiner, MD, has overseen recruitment of healthy controls, MCI and AD subjects from over 60 sites across the United States and Canada. (For up-to-date information, see [www.adni-info.org](http://www.adni-info.org)).

Ethical approval was obtained by the institutional review board at each participating center. All study participants provided written informed consent. Participants took part in the baseline and follow-up clinical, neuropsychometric, and MRI assessments. CSF data were collected from all subjects where possible.

Five diagnostic groups were used for this study: control, SMC, early MCI, late MCI, and AD. A diagnosis of MCI was based on a significant memory concern (by participant or informant) and a clinical dementia rating of 0.5, but a Mini-Mental State Examination no lower than 24. Level of MCI (early or late) was determined by the

Wechsler Memory Scale Logical Memory II. The SMC group was developed by ADNI to bridge the gap between controls and MCI participants, and is characterized by individuals who have self-reported a memory complaint but score within a normal range for cognition.

### 2.2. CSF measurements

CSF A $\beta$ 1-42, t-tau, and p-tau were measured at the ADNI biomarker core (University of Pennsylvania) using the microbead-based multiplex immunoassay, the INNO-BIA AlzBio3 RUO test (Fujirebio, Ghent, Belgium), on the Luminex platform (Luminex Corp, Austin, TX, USA). Raw, untransformed CSF values were used for all analyses.

Additional analyses were carried out on CSF data obtained using the Roche Elecsys cobas e 601 fully automated immunoassay platform at the ADNI biomarker core (University of Pennsylvania).

### 2.3. Images and image processing

MRI (3T) was acquired across sites using a standardized protocol that was validated across platforms and described in the study by Jack (2015). Images underwent quality control at the Mayo Clinic (Rochester, MN) which included assessments for protocol compliance and image quality, as well as checks for any significant neurological/radiological abnormalities. Baseline T1-weighted and FLAIR images from newly recruited individuals (i.e., with a first visit in ADNI2 or GO) were downloaded from the ADNI database.

Bayesian Model Selection, a fully automated lesion segmentation pipeline, was applied to the coregistered image pairs (T1 and FLAIR) (Sudre et al., 2015). Briefly, Bayesian Model Selection uses a Gaussian mixture model to jointly model healthy tissue and unexpected intensity observations for the different anatomical brain tissue. The number of Gaussian components required to appropriately model each tissue type is dynamically optimized using a split and merge strategy. After fitting the model, candidate hyperintense WMH voxels are selected and the formed connected components classified as lesion or artifacts in a postprocessing step based on anatomy rules. The resultant probability map of WMH was integrated to calculate the WMH volume. Only supratentorial WMH were included which were located in the white matter and subcortical gray matter. Total intracranial volumes (TIV) were calculated using the Geodesic information flows (GIF) label fusion framework, using T1-weighted images only (Cardoso et al., 2015).

All automated WMH segmentations were visually inspected by a single rater who had been trained to segment WMH manually. The postprocessing stage of the lesion classification was iteratively improved so as to limit the number of unusable segmentations and improve segmentations where large sections of lesion had been excluded. 22 cases were improved in this way and rechecked by a single rater, 21 of which were useable. Altogether, 12 regions were considered unusable of 944 regions in separate individuals. 3 further individuals were excluded because of lack of demographic/diagnostic information.

To calculate the parietal and nonparietal WMH, Euclidean distance maps from the cortical lobar parcellation obtained from GIF were used to assign each WM voxel to the closest associated lobe.

### 2.4. Statistical analyses

All statistical analysis was performed using STATA v13 (Stata Corp.). For WMH, most analyses were performed on log<sub>e</sub> transformed values to reduce skewness. To estimate differences in means across groups for demographic, imaging, and CSF variables, linear regression models were used for continuous variables. For

$\log_e$ WMH, linear regression models were used with group as the predictor of interest and adjustment for TIV. We also present untransformed WMH volumes and again test for differences in these across diagnostic groups, using linear regression with adjustment for TIV.

For analyses involving associations between WMH volume and CSF biomarkers, CSF biomarker values were standardized using the pooled within-group standard deviations (SDs) calculated from a linear regression model for each biomarker that allowed for differences in mean levels by group. Because WMH volumes were analyzed on a logarithmic scale the resulting coefficients were back-transformed so that they expressed percentage increases or decreases in WMH for each 1SD increase in CSF biomarkers.

To explore whether CSF A $\beta$ 1-42 was associated with WMH across diagnostic groups, separate diagnostic group-specific linear regression models were fitted with  $\log_e$ WMH as the dependent variable and standardized A $\beta$ 1-42 and TIV as predictors. Between-group differences in slope were assessed by fitting a further model in the combined diagnostic groups, with interactions between diagnostic group and A $\beta$ 1-42, and carrying out a joint interaction test. All models were also refitted with age at the baseline as an additional covariate. Analyses including age were secondary, since age at the baseline may be on the causal pathway (theoretically a proxy for the number of vascular events that may have happened to an individual and the ability to clear amyloid). Analogous procedures were used with A $\beta$ 1-42 values being replaced by t-tau and then by p-tau.

Scatter plots of  $\log_e$ WMH volume against CSF A $\beta$ 1-42, t-tau, or p-tau in each group were also generated to show unadjusted associations.

We decided that should the joint interaction terms between participant group and A $\beta$ 1-42 in the above models not be statistically significant then any further analysis would be carried out in the cohort as a whole, adjusting for participant group but without including the interaction terms. In a first set of linear regression models, we explored the A $\beta$ 1-42-WMH relationship adjusting for (1) nothing; (2) TIV; (3) TIV and diagnostic group; (4) TIV, group and age; (5) TIV, group age and APOE- $\epsilon$ 4 status. This was repeated in a second set of models adding t-tau as a covariate to all models in the set (1) to (5) and in a third set of models by adding p-tau to all models in the set (1) to (5). Partial  $R^2$  values for these potential confounders were also calculated for each covariate from model (5) in each set.

Supplementary analyses were also performed assessing the relationship between age and WMH using linear regression models with  $\log_e$ WMH being the dependent variable and age and TIV being predictors. This was initially performed in each diagnostic group separately, with one further model fitted that allowed for an interaction between diagnostic group and age. To assess the impact of vascular risk factors on the WMH-age relationship, smoking, hypertension, body mass index (BMI), and diabetes were added as covariates into separate diagnostic group-specific models and then in a combined model. To examine relationships between age and CSF variables, linear regression models were initially performed with A $\beta$ 1-42 as the dependent variable and age as the predictor, before replacing A $\beta$ 1-42 with t-tau and then p-tau. We also assessed associations between CSF biomarkers and parietal lobe WMH volumes versus nonparietal WMH. These analyses used CSF A $\beta$ 1-42, t-tau, or p-tau as the dependent variable and  $\log_e$  transformed parietal and nonparietal WMH volumes as predictor variables. The main analyses were repeated using the Elecsys biomarker data. These included the group-specific linear regression models with  $\log_e$ WMH as the dependent variable and standardized biomarker and TIV as predictors, and the analysis of CSF A $\beta$ 1-42 and WMH in the cohort as a whole.

### 3. Results

#### 3.1. Group demographics

There were 929 subjects with useable WMH values and available demographic data. [Table 1](#) shows demographic, imaging, and CSF biomarker summary statistics for each diagnostic group. Between-group differences were seen in age, with the AD group being the oldest and the EMCI group being the youngest. Participants differed as expected in terms of Mini-Mental State Examination, APOE- $\epsilon$ 4 status, WMH volume, CSF A $\beta$ 1-42, CSF t-tau, and CSF p-tau with poorer scores, greater  $\epsilon$ 4 carriage, and CSF biomarkers indicative of more pathology from controls to EMCI to LMCI to AD. In terms of vascular risk factors, differences were only observed between groups in BMI, with a lower BMI observed in the AD group. Summary CSF marker statistics for the subgroup with Elecsys CSF A $\beta$ 1-42 measures are shown in [Supplementary Table 1](#).

#### 3.2. CSF biomarkers and WMH

[Table 2](#) shows the estimated percentage decreases in WMH for increases in CSF A $\beta$ 1-42 after adjustment for TIV and [Fig. 1](#) shows the corresponding scatter plots. Although the estimated slope of the association in the 5 diagnostic groups varied somewhat (and was not formally statistically significant in the LMCI and AD groups), 95% confidence intervals were relatively wide and the variability in estimates was consistent with chance (joint tests for differences in slopes among groups were not statistically significant). The effect of CSF A $\beta$ 1-42 on WMH was typically reduced after adjustment for age, but we did not find any material changes in effects after adjustment for gender ([Supplementary Table 2](#)). [Supplementary Table 3](#) shows equivalent results to [Table 2](#) using the Elecsys CSF A $\beta$ 1-42 assay. These results demonstrate similar findings using this alternative assay.

[Table 3](#) shows the percentage increases in WMH for increases in CSF t-tau and [Fig. 2](#) shows the corresponding scatter plots. With adjustment for TIV, WMH burden was borderline positively associated with t-tau only in the EMCI group. This association was no longer significant after adjusting for age. There was no evidence for an interaction between t-tau and group.

[Table 4](#) shows the percentage increases in WMH for increases in CSF p-tau and [Fig. 3](#) shows the corresponding scatter plots. There was no evidence for an association between WMH and p-tau in any diagnostic group. There no evidence for an interaction between group and p-tau.

#### 3.3. Relationships of CSF A $\beta$ 1-42 with WMH

[Table 5a](#) shows the percentage reductions in WMH volume for a 1 SD increase in CSF A $\beta$ 1-42 adjusting for group, age, APOE- $\epsilon$ 4 status, t-tau, and p-tau. The relationship between WMH and A $\beta$ 1-42 remains significant and materially unchanged (between a 16% and 22% reduction in WMH per 1SD change in A $\beta$ 1-42) after progressive adjustment for covariates. This association between CSF A $\beta$ 1-42 and WMH was also shown to be independent of the effects of CSF t-tau and p-tau.

[Table 5b](#) shows partial  $R^2$  values for all the covariates in each model set. A $\beta$ 1-42 consistently uniquely explained approximately 5% of the variance in WMH across all models. Age explained the largest variance in WMH at ~15%, compared with tau biomarkers, group, and APOE- $\epsilon$ 4 status, with each only explaining only around 1%.

[Supplementary Table 4](#) shows analogous results to [Table 5](#) using the Elecsys assay. The results are not materially changed by the use of this alternative CSF A $\beta$ 1-42 measurement.

**Table 1**  
Demographic, imaging, and CSF summary statistics

Characteristic	Controls	SMC	EMCI	LMCI	AD	p Value across groups
N (total = 929)	180	107	320	171	151	
Age	73.4 (6.2)	72.3 (5.5)	71.0 (7.5)	72.4 (7.7)	75.0 (8.0)	<b>&lt;0.001</b>
Male (%)	46	43	54	56	56	0.08
MMSE/30	29.0 (1.3)	29.0 (1.3)	28.3 (1.6)	27.6 (1.8)	23.1 (2.1)	<b>&lt;0.001</b>
APOE-ε4 positive (%)	33	36	47	60	71	<b>&lt;0.001</b>
TIV (mL)	1403 (134)	1414 (125)	1423 (136)	1436 (132)	1416 (151)	0.2
WMH volume (mL)	3.4 (4.8)	3.4 (4.4)	3.8 (6.1)	3.7 (8.1)	5.8 (9.0)	<b>0.005<sup>a</sup></b>
Median (IQR)						
Log <sub>e</sub> WMH (mL)	1.3 (0.9)	1.3 (0.9)	1.4 (1.0)	1.4 (1.0)	1.7 (1.0)	<b>0.002<sup>a</sup></b>
CSF Aβ1-42 (pg/mL) <sup>b</sup>	270.5 (80.2)	308.3 (91.2)	251.0 (79.0)	212.0 (77.3)	181.1 (64.3)	<b>&lt;0.001</b>
Median (IQR)	287.5 (120.5)	318.0 (157.0)	255.0 (136.0)	190.5 (103.0)	169.0 (53.0)	
CSF t-tau (pg/mL) <sup>c</sup>	62.5 (29.0)	59.6 (24.7)	70.9 (42.3)	93.3 (50.2)	120.1 (56.3)	<b>&lt;0.001</b>
Median (IQR)	53.0 (33.6)	53.6 (27.3)	59.2 (39.1)	81.3 (72.6)	108.0 (71.2)	
CSF p-tau (pg/mL) <sup>d</sup>	21.8 (10.0)	21.6 (10.6)	23.0 (11.0)	28.6 (14.3)	34.2 (17.8)	<b>&lt;0.001</b>
Median (IQR)	19.6 (11.1)	18.3 (10.6)	20.4 (12.9)	25.5 (17.0)	31.3 (17.9)	
History of hypertension (%)	49	47	50	45	52	0.8
Smoking: never (%)	58	53	59	65	61	0.4
Previous (%)	37	45	39	33	34	
Current (%)	4	2	2	2	4	
BMI	27.4 (4.6)	28.4 (6.3)	28.2 (5.3)	27.4 (4.9)	26.2 (5.3)	<b>&lt;0.001</b>
History of diabetes (%)	7	10	11	9	15	0.2

Values are mean (SD) unless reported. Bold values denote statistical significance at the  $p < 0.05$  level.

Key: SMC, significant memory concern; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD, Alzheimer's disease; MMSE, Mini Mental State Examination; TIV, total intracranial volume; WMH, white matter hyperintensities; BMI, body mass index.

<sup>a</sup> Adjusted for TIV.

<sup>b</sup> Available in 144 controls, 92 SMC, 265 EMCI, 150 LMCI, 125 AD.

<sup>c</sup> Available in 139 controls, 92 SMC, 252 EMCI, 145 LMCI, 119 AD.

<sup>d</sup> Available in 143 controls, 92 SMC, 265 EMCI, 149 LMCI, 125 AD.

#### 4. Discussion

In this study, we found strong evidence for a consistent association between CSF Aβ1-42 and WMH in all diagnostic groups, whereby a decrease in CSF Aβ1-42 was associated with an increase in WMH, that was independent of CSF t-tau and independent of CSF p-tau. Furthermore, this relationship remained significant in the pooled cohort when adjusting for age, diagnostic group, TIV, and APOE-ε4 status. There was very little evidence for a relationship between t-tau pathology and WMH, with a borderline positive association only in the EMCI group. No association of p-tau with WMH was observed. To our knowledge, this is the first study to look for a direct association between Aβ1-42 and tau CSF biomarkers and WMH volumes across the disease spectrum, from normal aging to AD, including patients with EMCI and SMC.

Our findings provide evidence for a link between WMH and amyloid pathology, from normal aging through to clinical AD. Notably, evidence from autosomal dominantly inherited familial AD suggests that changes in WMH could occur in the presymptomatic

stage of disease, deviating from normal controls around 6 years before symptom onset (Lee et al., 2016). Our observations from groups at higher risk of developing sporadic AD dementia add further weight to the notion that WMH are a core feature of AD.

We did not find any evidence for differing relationships in WMH and Aβ1-42, p-tau, or t-tau between diagnostic groups. WMH are heterogeneous and are likely to represent multiple pathologies that could potentially vary throughout the disease course. Given the recent evidence that WMH may be more related to neurodegenerative and tau pathologies in clinical AD (McAleese et al., 2017), it would have been reasonable to propose stronger relationships with tau at later disease stages. Because Aβ1-42 levels are thought to be relatively stable in clinical AD stages (Zhou et al., 2009), it could also be expected that there might be less of an association with WMH in later disease stages. The fact that we did not find any differences between disease stages suggests that WMH relationships with AD CSF biomarkers actually remain consistent across the disease course.


Age at the baseline was a stronger independent predictor of WMH than Aβ1-42, which likely reflects the contributions of

**Table 2**  
Regression model results for the relationship between WMH and CSF Aβ1-42, by group

WMH and Aβ1-42	Controls	SMC	EMCI	LMCI	AD
TIV adjusted					
% reduction in WMH (mL) per 1SD change in Aβ1-42	30.4	20.0	22.0	14.9	18.0
95% Confidence intervals	(20.3–39.3)	(6.4–31.6)	(12.5–30.5)	(-0.5–28.0)	(-0.2–32.8)
p Value	<b>&lt;0.001</b>	<b>0.006</b>	<b>&lt;0.001</b>	0.06	0.05
TIV and age adjusted					
% reduction in WMH (mL) per 1SD change in Aβ1-42	26.7	18.7	12.3	9.4	21.7
95% Confidence intervals	(15.7–36.2)	(5.2–30.2)	(2.5–21.0)	(-5.5–22.2)	(6.8–34.2)
p Value	<b>&lt;0.001</b>	<b>0.009</b>	<b>0.02</b>	0.2	<b>0.006</b>
Between group differences in % reductions per 1SD change in Aβ1-42 : p value = 0.4 (0.4 with age adjustment)					

Estimates are percentage decrease in WMH volume per one SD change (1 SD = 78.3 pg/mL) in Aβ1-42. The SD is the pooled within-group SD, calculated from a linear regression model that allowed for differences in mean levels of Aβ1-42 by group. Bold values denote statistical significance at the  $p < 0.05$  level.

Key: SMC, significant memory concern; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD, Alzheimer's disease; TIV, total intracranial volume; WMH, white matter hyperintensities.



**Fig. 1.** Scatter plots showing the relationship between WMH and CSF A $\beta$ 1-42 in controls, significant memory concern (SMC), early mild cognitive impairment (EMCI), late mild cognitive impairment (LMCI), Alzheimer's disease (AD) groups. WMH volume is shown on a log scale and is unadjusted for any covariates. Abbreviations: WMH, white matter hyperintensities.

vascular risk factors, vascular events, and comorbidities that accrue with age. Taken together, although our study is purely correlative, our results are compatible with the theory that some WMH are part of the AD pathological process, and some are the result of an aging/vascular process (McAleese et al., 2015, 2017). Although we did not find any evidence for contributions from smoking, hypertension, BMI, and diabetes to the relationship between WMH and age (Supplementary Table 5), a more comprehensive examination of the effects of vascular risk factors, vascular disease, and vascular-related events on the age-WMH relationship may provide greater insight into the potential causes of WMH associated with age.

In studies of A $\beta$  and WMH, there have been mixed findings; some have reported a lack of any association (Kalheim et al., 2017; Kester et al., 2014), while several studies confirm our findings that lower CSF A $\beta$ 1-42 is associated with higher WMH in both controls and disease groups (Marnane et al., 2016; Pietrobboni et al., 2018). Some have reported that lower levels of CSF A $\beta$ 1-42 were particularly associated with WMH in AD subjects, as opposed to controls or MCI subjects (Shams et al., 2016; Van Westen et al., 2016). Results from amyloid PET studies have also demonstrated significant associations with WMH in controls (Kandel et al., 2016; Marnane et al., 2016), MCI and AD groups (Zhou et al., 2015). Although a


**Table 3**

Regression model results for the relationship between WMH and CSF t-tau, by group

WMH and t-tau	Controls	SMC	EMCI	LMCI	AD
TIV adjusted					
% Increase in WMH (mL) per 1SD change in t-tau	5.1	3.8	13.2	-4.0	-3.3
95% Confidence intervals	(-16.4 to 32.0)	(-24.9 to 43.5)	(0.2 to 27.7)	(-16.9 to 11.1)	(-15.2 to 10.2)
p Value	0.7	0.8	0.05	0.6	0.6
TIV and age adjusted					
% Increase in WMH (mL) per 1SD change in t-tau	-7.3	-5.1	2.4	-3.9	3.6
95% Confidence intervals	(-26.2 to 16.3)	(-31.2 to 30.9)	(-8.0 to 14.0)	(-15.7 to 9.5)	(-7.9 to 16.5)
p Value	0.5	0.8	0.7	0.6	0.6
Between group differences in % increases per 1SD change in t-tau : p value = 0.4 (0.7 with age adjustment)					

Estimates are percentage increase in WMH volume per one SD change (1 SD = 42.8 pg/mL) in t-tau. The SD is the pooled within-group SD, calculated from a linear regression model that allowed for differences in mean levels of t-tau by group.

Key: SMC, significant memory concern; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD, Alzheimer's disease; TIV, total intracranial volume; WMH, white matter hyperintensities.



**Fig. 2.** Scatter plots showing the relationship between WMH and CSF t-tau in controls, significant memory concern (SMC), early mild cognitive impairment (EMCI), late mild cognitive impairment (LMCI), and Alzheimer's disease (AD) groups. WMH volume is shown on a log scale and is unadjusted for any covariates. Abbreviation: WMH, white matter hyperintensities.

recent systematic review concluded that most studies demonstrate a lack of significant relationship between amyloid PET and WMH burden (Roseborough et al., 2017).

This study adds to the expanding body of literature attempting to characterize the complex relationship between AD and presumed vascular pathology at different disease stages. As well as evidence that both A $\beta$  deposition and WMH independently predict cognitive impairment or brain atrophy (Barnes et al., 2013; Gordon et al., 2015; Haight et al., 2013; Provenzano et al., 2013), a growing number of

studies are suggestive of an additive effect of combined pathology on neurodegeneration (Bos et al., 2018) and/or cognitive decline (Lopez et al., 2014; Vemuri et al., 2015; Vemuri and Knopman, 2016). Several studies have investigated the possibility of an interactive effect between the 2 pathologies, whereby there is a direct influence of one pathology on the other. One study found that WMH predicted hippocampal atrophy in preclinical AD subjects with abnormal CSF A $\beta$  (Freeze et al., 2016), and another found that low CSF A $\beta$  predicted greater WMH accrual over time (Scott et al., 2016).


**Table 4**

Regression model results for the relationship between WMH and CSF p-tau, by group

WMH and p-tau	Controls	SMC	EMCI	LMCI	AD
TIV adjusted					
% Increase in WMH (mL) per 1SD change in p-tau	13.6	14.1	3.5	-8.7	0.5
95% Confidence intervals	(-6.6 to 38.3)	(-8.8 to 42.8)	(-10.0 to 18.9)	(-21.3 to 5.9)	(-10.8 to 13.3)
p Value	0.2	0.3	0.6	0.2	0.9
TIV and age adjusted					
% Increase in WMH (mL) per 1SD change in p-tau	7.3	13.1	-3.9	-3.0	8.0
95% Confidence intervals	(-11.3 to 29.9)	(-9.0 to 40.5)	(-14.8 to 8.4)	(-15.3 to 11.1)	(-3.0 to 20.3)
p Value	0.5	0.3	0.5	0.7	0.1
Between group differences in % increases per 1SD change in p-tau : p Value = 0.4 (0.5 with age adjustment)					

Estimates are percentage increase in WMH volume per one SD (1SD = 12.8 pg/mL) change in p-tau. The SD is the pooled within-group SD, calculated from a linear regression model that allowed for differences in mean levels of p-tau by group.

Key: SMC, significant memory concern; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD, Alzheimer's disease; TIV, total intracranial volume; WMH, white matter hyperintensities.



**Fig. 3.** Scatter plots showing the relationship between WMH and CSF p-tau in controls, significant memory concern (SMC), early mild cognitive impairment (EMCI), late mild cognitive impairment (LMCI), and Alzheimer's disease (AD) groups. WMH volume is shown on a log scale and is unadjusted for any covariates. Abbreviation: WMH, white matter hyperintensities.

Hypotheses into mechanistic links between A $\beta$  and WMH have mainly focused on the proposed vascular origin of WMH and reflect the idea that there may be positive feedback between the 2 pathologies (Smith and Greenberg, 2009; Zlokovic, 2011). Cerebral amyloid angiopathy (CAA) may be a potential mediator of the relationship between WMH and A $\beta$ , whereby SVD could lead to reduced clearance of A $\beta$  from brain parenchyma, enhancing its deposition in cerebral vessel walls. This in turn may result in cerebrovascular dysfunction, leading to eventual hypoperfusion and ischemia and subsequent white matter lesions that appear as WMH (Grinberg and Thal, 2010; Hawkes et al., 2014; Weller et al., 2008). Previous research has demonstrated associations of CAA with WMH (Ryan et al., 2015) and with CSF A $\beta$  (Renard et al., 2012; Verbeek et al., 2009). A recent paper has shown that WMH related to A $\beta$  are associated with cerebral microbleeds (a surrogate of CAA pathology) (Graff-Radford et al., 2019). In addition to the vascular origin of WMH, recent evidence suggests that parietal WMH could be a result of degenerative axonal loss caused by AD pathology (McAleese et al., 2015, 2017). In this study though, we did not find any evidence for a stronger association between CSF biomarkers and WMH in the parietal lobe (Supplementary Table 6).

There are comparatively fewer studies regarding the association with tau and WMH and results are often conflicting. A positive association has been demonstrated neuropathologically (McAleese et al., 2015) and with PET (Vemuri et al., 2017); more direct measures of brain tau load than CSF measures. A positive association has been demonstrated over time in CSF, but the relationship is less

clear with baseline measures of tau (Tosto et al., 2015). Our study did not find much evidence of a relationship between tau pathology and WMH, which is in line with several other studies (Guzman et al., 2013; Kester et al., 2014; Osborn et al., 2018).

The fact that age was the strongest predictor of WMH is consistent with previous studies (de Leeuw et al., 2001; Raz et al., 2012; Schneider et al., 2009). We also found that the association between biomarkers and age was not uniform between diagnostic groups (Supplementary Table 7). In controls and EMCI groups, older ages were associated with lower A $\beta$ 1-42 and higher tau, but in the AD group, younger ages were associated with higher tau. This differential effect of age on CSF biomarkers between AD groups and controls has been previously reported (Mattsson et al., 2012).

Investigations of the associations between AD CSF biomarkers and WMH in previous studies have either been performed on small samples or in studies with variable acquisition and analysis protocols. (Bos et al., 2018; Freeze et al., 2016; Kalheim et al., 2017; Pietroboni et al., 2018; Tosto et al., 2015). Using the ADNI cohort therefore provides the benefits of a multicenter standardized data acquisition protocol, as well as participants who have been systematically evaluated. Our study adds to previous findings using the ADNI cohort (Guzman et al., 2013; Haight et al., 2013; Marnane et al., 2016; Scott et al., 2016), in that data from all diagnostic groups are used and biomarkers for both tau and A $\beta$  are compared with quantitative WMH. We also treated both our biomarker and WMH data as continuous variables; most previous studies have dichotomized variables.

**Table 5**  
Regression model results examining the association between CSF A $\beta$ 1–42 and WMH with various covariates

(a)				
Model set	Covariates	% Reduction in WMH per 1SD change in A $\beta$ 1–42	95% Confidence interval	p Value
1 n = 776	i) None	21.9	17.1 to 26.5	<0.001
	ii) TIV	21.5	16.7 to 26.0	<0.001
	iii) TIV, group	21.6	16.2 to 26.6	<0.001
	iv) TIV, group, age	16.4	11.2 to 21.3	<0.001
	v) TIV, group, age, APOE-e4 status	17.9	12.2 to 23.2	<0.001
2 n = 747	i) T-tau	24.5	19.2 to 29.5	<0.001
	ii) T-tau, TIV	23.2	17.9 to 28.1	<0.001
	iii) T-tau, TIV, group	23.1	17.5 to 28.3	<0.001
	iv) T-tau, TIV, group, age	18.0	12.6 to 23.1	<0.001
	v) T-tau, TIV, group, age, APOE-e4 status	18.8	13.0 to 24.2	<0.001
3 n = 774	i) P-tau	24.1	18.7 to 29.1	<0.001
	ii) P-tau, TIV	23.8	18.6 to 28.7	<0.001
	iii) P-tau, TIV, group	23.8	18.2 to 29.0	<0.001
	iv) P-tau, TIV, group, age	17.7	12.2 to 23.0	<0.001
	v) P-Tau, TIV, group, age, APOE-e4 status	18.9	12.9 to 24.4	<0.001

(b)		
Model set	Covariates from v)	Partial R <sup>2</sup>
1	A $\beta$ 1–42	0.05
	TIV	0.05
	Group	0.003
	Age	0.16
	APOE-e4 status	0.002
2	A $\beta$ 1–42	0.05
	t-tau	0.005
	TIV	0.06
	Group	0.005
	Age	0.16
3	APOE-e4 status	0.0007
	A $\beta$ 1–42	0.05
	p-tau	0.003
	TIV	0.06
	Group	0.004
	Age	0.15
	APOE-e4 status	0.001

All models assess associations between WMH and A $\beta$ 1–42 with various covariates, with model 2 additionally adjusting for t-tau and model 3 additionally adjusting for p-tau. In a) estimates are percentage changes in WMH volume per one SD change (1 SD = 78.3 pg/mL) in A $\beta$ 1–42. The SD is the pooled within-group SD, calculated from a linear regression model that allowed for differences in mean levels of A $\beta$ 1–42 by group. In b) partial R<sup>2</sup> are shown for each covariate in the fully-adjusted model v) from each of the 3 model sets. R<sup>2</sup> values are shown to 2 significant figures for values >0.1 and 1 significant figure for values <0.1.

Key: TIV, total intracranial volume; WMH, white matter hyperintensities.

There are several limitations to this work, which include the selection criteria for the ADNI cohort meaning any generalizations to the wider population must be made cautiously. ADNI participants tend to be of high socioeconomic status, well-educated and, importantly for this study, free of large amounts of cerebrovascular disease (a score of 4 or lower on the Hachinski Ischemic Scale) (<https://adni.loni.usc.edu/wp-content/uploads/2008/07/adni2-procedures-manual.pdf>). This study was also only carried out using baseline data; longitudinal studies are necessary to characterize the relationship between WMH and AD biomarkers over time, an approach which will be facilitated by the increasing use of volumetric FLAIR imaging providing more precise measures of change.

There are limitations surrounding the use of CSF biomarkers in this study to investigate AD pathology. CSF biomarkers can now be measured by several different assays, introducing the possibility

that these methodological differences may influence study findings. However, [Supplementary Tables 3 and 4](#) demonstrate that analyses carried out using another assay platform, Elecsys, confirm our findings of the associations between CSF biomarkers and WMH. In addition, despite the benefits of using continuous biomarker variables, it would be interesting to look at associations within biomarker-negative and biomarker-positive individuals to explore whether associations may be driven by those on the amyloid pathway ([Jack et al., 2018](#)) or those who are not. CSF biomarkers are increasingly used to provide information about underlying biochemical changes associated with disease, with A $\beta$ 1–42 and p-tau well established as markers of AD pathology and t-tau as an established marker of neurodegeneration ([Blennow et al., 2010; Jack et al., 2018](#)). However, measures of actual brain A $\beta$  and tau burden and their local distribution, through PET imaging or pathological investigations, were either not available or not used.

This study only examined associations between WMH and the well-established AD biomarkers of CSF A $\beta$ 1–42, t-tau, and p-tau. The existence of additional copathologies in AD suggest that further biomarkers are needed to complement these classical measurements. One biomarker of particular relevance is neurofilament light, a marker of large caliber axonal injury ([Skillback et al., 2014](#)). Previous studies of associations between WMH and neurofilament light have reported mixed results ([Mattsson et al., 2019; Osborn et al., 2018; Zetterberg et al., 2016](#)), and therefore future analysis across the full spectrum of the disease course using our well-characterized WMH segmentation tool would be highly pertinent.

A lack of pathological confirmation of cases in the ADNI cohort may, in part, explain our findings. Diagnostic groups may be bimodal in nature; control groups may contain atypical or early stage AD subjects along with the cognitively normal, and a significant proportion of MCI subjects supposedly at high risk for conversion to AD will not have any underlying AD pathology ([Jicha et al., 2006; Petersen et al., 2006](#)). Indeed in our study, this may have been a contributing factor to the lack of observed differences in biomarker and WMH relationships between diagnostic groups.

In addition, although the 4 measures of vascular risk used in this study are unlikely to fully characterize all aspects of SVD, they are among the most notable risk factors, as well as being the most consistently documented for the ADNI cohort.

## 5. Conclusion

In summary, our study suggests that CSF A $\beta$ 1–42 is a consistent predictor of WMH across all diagnostic groups, in a manner that is not dependent on CSF tau. As WMH are thought to be largely vascular in origin, this study adds to the literature exploring the complex relationship between amyloid and presumed vascular pathologies. Understanding the different relationships between disease markers are important to build more realistic models of biomarkers changes across the full spectrum of normal aging to AD.

## Disclosure statement

The authors have no actual or potential conflicts of interest.

## CRedit authorship contribution statement

**Phoebe Walsh:** Conceptualization, Data curation, Methodology, Writing - original draft, Writing - review & editing. **Carole H. Sudre:** Methodology, Software. **Cassidy M. Fiford:** Methodology, Data curation. **Natalie S. Ryan:** Supervision. **Tammarny Lashley:** Supervision. **Chris Frost:** Methodology, Formal analysis, Writing - review & editing. **Josephine Barnes:** Conceptualization, Writing - review & editing, Supervision.



## Acknowledgements

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc; Cogstate; Eisai Inc; Elan Pharmaceuticals, Inc; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc; Fujirebio; GE Healthcare; IXICO Ltd; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co, Inc; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health ([www.fnih.org](http://www.fnih.org)). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

P Walsh is supported by the Wolfson Foundation. J Barnes and C Fiford were funded by an Alzheimer's Research UK Senior Fellowship. C Sudre is supported by an Alzheimer's Society Junior Fellowship (AS-JF-17-011). N Ryan is supported by a University of London Chadburn Academic Clinical Lectureship. T Lashley is supported by an Alzheimer's Research UK senior fellowship. The Dementia Research Centre is supported by Alzheimer's Research UK, Brain Research Trust, and The Wolfson Foundation. This work was supported by the NIHR Queen Square Dementia Biomedical Research Unit and the National Institute for Health Research Biomedical Research Centre (BRC).

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.neurobiolaging.2020.03.008>.

## References

- Barnes, J., Carmichael, O.T., Leung, K.K., Schwarz, C., Ridgway, G.R., Bartlett, J.W., Malone, I.B., Schott, J.M., Rossor, M.N., Biessels, G.J., Decarli, C., Fox, N.C., Alzheimer's Disease Neuroimaging, I., 2013. Vascular and Alzheimer's disease markers independently predict brain atrophy rate in Alzheimer's Disease Neuroimaging Initiative controls. *Neurobiol. Aging* 34, 1996–2002.
- Blennow, K., Hampel, H., Weiner, M., Zetterberg, H., 2010. Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat. Rev. Neurol.* 6, 131–144.
- Bos, I., Verhey, F.R., Ramakers, I.H.G.B., Jacobs, H.I.L., Soininen, H., Freund-Levi, Y., Hampel, H., Tsolaki, M., Wallin, A.K., Van Buchem, M.A., Oleksik, A., Verbeek, M.M., Rikkert, M.O., Van Der Flier, W.M., Scheltens, P., Aalten, P., Visser, P.J., Vos, S.J.B., 2018. Correction to: cerebrovascular and amyloid pathology in pre-dementia stages: the relationship with neurodegeneration and cognitive decline (Alzheimer's Res. Ther. (2017) 9 DOI: 10.1186/s13195-017-0328-9). *Alzheimer's Res. Ther.* 10, 1–10.
- Brayne, C., Richardson, K., Matthews, F.E., Fleming, J., Hunter, S., Xuereb, J.H., Paykel, E., Mukaetova-Ladinska, E.B., Huppert, F.A., O'Sullivan, A., Denning, T., Cambridge City Over-75s Cohort Cc75c Study Neuropathology, C., 2009. Neuropathological correlates of dementia in over-80-year-old brain donors from the population-based Cambridge city over-75s cohort (CC75C) study. *J. Alzheimers Dis.* 18, 645–658.
- Cardoso, M.J., Modat, M., Wolz, R., Melbourne, A., Cash, D., Rueckert, D., Ourselin, S., 2015. Geodesic information flows: Spatially-Variant Graphs and their Application to segmentation and fusion. *IEEE Trans. Med. Imaging* 34, 1976–1988.
- De Leeuw, F.E., DE Groot, J.C., Van Gijn, J., 2001. [Cerebral white matter lesions in the elderly: vascular risk factors and cognitive consequences]. *Ned Tijdschr Geneesk* 145, 2067–2071.
- Esiri, M.M., Joachim, C., Sloan, C., Christie, S., Agacinski, G., Bridges, L.R., Wilcock, G.K., Smith, A.D., 2014. Cerebral subcortical small vessel disease in subjects with pathologically confirmed Alzheimer disease: a clinicopathologic study in the Oxford Project to Investigate Memory and Ageing (OPTIMA). *Alzheimer Dis. Assoc. Disord.* 28, 30–35.
- Freeze, W.M., Jacobs, H.I.L., Gronenschild, E.H., Jansen, J.F.A., Burgmans, S., Aalten, P., Clerx, L., Vos, S.J., VAN Buchem, M.A., Barkhof, F., Van Der Flier, W.M., Verbeek, M.M., Rikkert, M.O., Backes, W.H., Verhey, F.R., 2016. White matter hyperintensities potentiate hippocampal volume reduction in non-demented older individuals with abnormal amyloid- $\beta$ . *J. Alzheimer's Dis.* 55, 333–342.
- Gordon, B.A., Najmi, S., Hsu, P., Roe, C.M., Morris, J.C., Benzinger, T.L.S., 2015. The effects of white matter hyperintensities and amyloid deposition on Alzheimer dementia. *NeuroImage Clin.* 8, 246–252.
- Gorelick, P.B., Scuteri, A., Black, S.E., Decarli, C., Greenberg, S.M., Iadecola, C., Launer, L.J., Laurent, S., Lopez, O.L., Nyenhuis, D., Petersen, R.C., Schneider, J.A., Tzourio, C., Arnett, D.K., Bennett, D.A., Chui, H.C., Higashida, R.T., Lindquist, R., Nilsson, P.M., Roman, G.C., Sellke, F.W., Seshadri, S., 2011. Vascular contributions to cognitive impairment and dementia: a statement for healthcare professionals from the American heart association/American stroke association. *Stroke* 42, 2672–2713.
- Gouw, A.A., Seewann, A., van der Flier, W.M., Barkhof, F., Rozemuller, A.M., Scheltens, P., Geurts, J.J., 2011. Heterogeneity of small vessel disease: a systematic review of MRI and histopathology correlations. *J. Neurol. Neurosurg. Psychiatry* 82, 126–135.
- Graff-Radford, J., Arenaza-Urquijo, E.M., Knopman, D.S., Schwarz, C.G., Brown Jr., R.D., Rabinstein, A.A., Gunter, J.L., Senjem, M.L., Przybelski, S.A., Lesnick, T., Ward, C., Mielke, M.M., Lowe, V.J., Petersen, R.C., Kremers, W.K., Kantarci, K., Jack Jr., C.R., Vemuri, P., 2019. White matter hyperintensities: relationship to amyloid and tau burden. *Brain*.
- Grinberg, L.T., Thal, D.R., 2010. Vascular pathology in the aged human brain. *Acta Neuropathologica* 119, 277–290.
- Guzman, V.A., Carmichael, O.T., Schwarz, C., Tosto, G., Zimmerman, M.E., Brickman, A.M., 2013. White matter hyperintensities and amyloid are independently associated with entorhinal cortex volume among individuals with mild cognitive impairment. *Alzheimer's Dement.* 9, S124–S131.
- Haight, T.J., Landau, S.M., Carmichael, O., Schwarz, C., Decarli, C., Jagust, W.J., 2013. Dissociable effects of Alzheimer disease and white matter hyperintensities on brain metabolism. *JAMA Neurol.* 70, 1039–1045.
- Hawkes, C.A., Jayakody, N., Johnston, D.A., Bechmann, I., Carare, R.O., 2014. Failure of perivascular drainage of  $\beta$ -amyloid in cerebral amyloid angiopathy. *Brain Pathol.* 24, 396–403.
- Jack, C.R., Bennett, D.A., Blennow, K., Carrillo, M.C., Dunn, B., Haeberlein, S.B., Holtzman, D.M., Jagust, W., Jessen, F., Karlawish, J., Liu, E., Molinuevo, J.L., Montine, T., Phelps, C., Rankin, K.P., Rowe, C.C., Scheltens, P., Siemers, E., Snyder, H.M., Sperling, R., Elliott, C., Masliah, E., Ryan, L., Silverberg, N., 2018. NIA-AA Research Framework: toward a biological definition of Alzheimer's disease. *Alzheimer's Dement.* 14, 535–562.
- Jack Jr., C.R., Barnes, J., Bernstein, M.A., Borowski, B.J., Brewer, J., Clegg, S., Dale, A.M., Carmichael, O., Ching, C., Decarli, C., Desikan, R.S., Fennema-Notestane, C., Fjell, A.M., Fletcher, E., Fox, N.C., Gunter, J., Gutman, B.A., Holland, D., Hua, X., Insel, P., Kantarci, K., Killiany, R.J., Krueger, G., Leung, K.K., Mackin, S., Maillard, P., Malone, I.B., Mattsson, N., Mcevoy, L., Modat, M., Mueller, S., Nosheny, R., Ourselin, S., Schuff, N., Senjem, M.L., Simonson, A., Thompson, P.M., Rettmann, D., Vemuri, P., Walhovd, K., Zhao, Y., Zink, S., Weiner, M., 2015. Magnetic resonance imaging in Alzheimer's Disease Neuroimaging Initiative 2. *Alzheimer's Dement.* 11, 740–756.
- Jellinger, K.A., Attems, J., 2006. Prevalence and impact of cerebrovascular pathology in Alzheimer's disease and parkinsonism. *Acta Neurol. Scand.* 114, 38–46.
- Jicha, G.A., Parisi, J.E., Dickson, D.W., Johnson, K., Cha, R., Ivnik, R.J., Tangalos, E.G., Boeve, B.F., Knopman, D.S., Braak, H., Petersen, R.C., 2006. Neuropathologic outcome of mild cognitive impairment following progression to clinical dementia. *Arch. Neurol.* 63, 674–681.
- Kalheim, L.F., Bjørnerud, A., Fladby, T., Vegge, K., Selnes, P., 2017. White matter hyperintensity microstructure in amyloid dysmetabolism. *J. Cereb. Blood Flow Metab.* 37, 356–365.
- Kandel, B.M., Avants, B.B., Gee, J.C., Mcmillan, C.T., Erus, G., Doshi, J., Davatzikos, C., Wolk, D.A., 2016. White matter hyperintensities are more highly associated with preclinical Alzheimer's disease than imaging and cognitive markers of neurodegeneration. *Alzheimer's Dement. Diagn. Assess. Dis. Monit.* 4, 18–27.
- Kapasi, A., Decarli, C., Schneider, J.A., 2017. Impact of multiple pathologies on the threshold for clinically overt dementia. *Acta Neuropathol.* 134, 171–186.
- Kester, M.I., Goos, J.D.C., Teunissen, C.E., Benedictus, M.R., Bouwman, F.H., Wattjes, M.P., Barkhof, F., Scheltens, P., van der Flier, W.M., 2014. Associations between cerebral small-vessel disease and Alzheimer disease pathology as measured by cerebrospinal fluid biomarkers. *JAMA Neurol.* 71, 855–862.
- Lee, S., Viqar, F., Zimmerman, M.E., Narkhede, A., Tosto, G., Benzinger, T.L.S., Marcus, D.S., Fagan, A.M., Goate, A., Fox, N.C., Cairns, N.J., Holtzman, D.M., Buckles, V., Ghetti, B., Mcdade, E., Martins, R.N., Saykin, A.J., Masters, C.L., Ringman, J.M., Ryan, N.S., Förster, S., Laske, C., Schofield, P.R., Sperling, R.A., Salloway, S., Correia, S., Jack, C., Weiner, M., Bateman, R.J., Morris, J.C., Mayeux, R., Brickman, A.M., 2016. White matter hyperintensities are a core feature of Alzheimer's disease: evidence from the dominantly inherited Alzheimer network. *Ann. Neurol.* 79, 929–939.

- Lopez, O.L., Klunk, W.E., Mathis, C., Becker, J.T., Aizenstein, H.J., Snitz, B., Cohen, A., Ikonomic, M., McDade, E., Dekosky, S.T., Weissfeld, L., Kuller, L., 2014. Amyloid, neurodegeneration, and small vessel disease as predictors of dementia in the oldest-old. *Neurology* 83, 1804–1811.
- Marnane, M., Al-Jawadi, O.O., Mortazavi, S., Pogorzelec, K.J., Wang, B.W., Feldman, H.H., Hsiung, G.Y.R., 2016. Periventricular hyperintensities are associated with elevated cerebral amyloid. *Neurology* 86, 535–543.
- Mattsson, N., Cullen, N.C., Andreasson, U., Zetterberg, H., Blennow, K., 2019. Association between longitudinal plasma neurofilament light and neurodegeneration in patients with Alzheimer disease. *JAMA Neurol.* 76, 791–799.
- Mattsson, N., Rosen, E., Hansson, O., Andreasen, N., Parnetti, L., Jonsson, M., Herukka, S.K., van der Flier, W.M., Blankenstein, M.A., Ewers, M., Rich, K., Kaiser, E., Verbeek, M.M., Olde Rikkert, M., Tsolaki, M., Mulugeta, E., Aarsland, D., Visser, P.J., Schroder, J., Marcusson, J., de Leon, M., Hampel, H., Scheltens, P., Wallin, A., Eriksdotter-Jonhagen, M., Minthon, L., Winblad, B., Blennow, K., Zetterberg, H., 2012. Age and diagnostic performance of Alzheimer disease CSF biomarkers. *Neurology* 78, 468–476.
- McAleese, K.E., Firbank, M., Dey, M., Colloby, S.J., Walker, L., Johnson, M., Beverley, J.R., Taylor, J.P., Thomas, A.J., O'Brien, J.T., Attems, J., 2015. Cortical tau load is associated with white matter hyperintensities. *Acta Neuropathologica Commun.* 3, 60.
- McAleese, K.E., Walker, L., Graham, S., Moya, E.L.J., Johnson, M., Erskine, D., Colloby, S.J., Dey, M., Martin-Ruiz, C., Taylor, J.P., Thomas, A.J., Mckeith, I.G., DE Carli, C., Attems, J., 2017. Parietal white matter lesions in Alzheimer's disease are associated with cortical neurodegenerative pathology, but not with small vessel disease. *Acta Neuropathol.* 134, 459–473.
- Osborn, K.E., Liu, D., Samuels, L.R., Moore, E.E., Cambroner, F.E., Acosta, L.M.Y., Bell, S.P., Babicz, M.A., Gordon, E.A., Pechman, K.R., Davis, L.T., Gifford, K.A., Hohman, T.J., Blennow, K., Zetterberg, H., Jefferson, A.L., 2018. Cerebrospinal fluid beta-amyloid42 and neurofilament light relate to white matter hyperintensities. *Neurobiol. Aging* 68, 18–25.
- Petersen, R.C., Parisi, J.E., Dickson, D.W., Johnson, K.A., Knopman, D.S., Boeve, B.F., Jicha, G.A., Ivnik, R.J., Smith, G.E., Tangalos, E.G., Braak, H., Kokmen, E., 2006. Neuropathologic features of amnesic mild cognitive impairment. *Arch. Neurol.* 63, 665–672.
- Pietroboni, A.M., Scarioni, M., Carandini, T., Basilico, P., Cadioli, M., Giulietti, G., Arighi, A., Caprioli, M., Serra, L., Sina, C., Fenoglio, C., Ghezzi, L., Fumagalli, G.G., de Riz, M.A., Calvi, A., Triulzi, F., Bozzali, M., Scarpini, E., Galimberti, D., 2018. CSF  $\beta$ -amyloid and white matter damage: a new perspective on Alzheimer's disease. *J. Neurol. Neurosurg. Psychiatry* 89, 352–357.
- Provenzano, F.A., Muraskin, J., Tosto, G., Narkhede, A., Wasserman, B.T., Griffith, E.Y., Guzman, V.A., Meier, I.B., Zimmerman, M.E., Brickman, A.M., 2013. White matter hyperintensities and cerebral amyloidosis: necessary and sufficient for clinical expression of Alzheimer disease? *JAMA Neurol.* 70, 455–461.
- Raz, N., Yang, Y., Dahle, C.L., Land, S., 2012. Volume of white matter hyperintensities in healthy adults: contribution of age, vascular risk factors, and inflammation-related genetic variants. *Biochim. Biophys. Acta* 1822, 361–369.
- Renard, D., Castelnuovo, G., Wacongne, A., Le Floch, A., Thouvenot, E., Mas, J., Gabelle, A., Labauge, P., Lehmann, S., 2012. Interest of CSF biomarker analysis in possible cerebral amyloid angiopathy cases defined by the modified Boston criteria. *J. Neurol.* 259, 2429–2433.
- Roseborough, A., Ramirez, J., Black, S.E., Edwards, J.D., 2017. Associations between amyloid  $\beta$  and white matter hyperintensities: a systematic review. *Alzheimer's Dement.* 13, 1154–1167.
- Ryan, N.S., Biessels, G.J., Kim, L., Nicholas, J.M., Barber, P.A., Walsh, P., Gami, P., Morris, H.R., Bastos-Leite, A.J., Schott, J.M., Beck, J., Mead, S., Chavez-Gutierrez, L., de Strooper, B., Rossor, M.N., Revesz, T., Lashley, T., Fox, N.C., 2015. Genetic determinants of white matter hyperintensities and amyloid angiopathy in familial Alzheimer's disease. *Neurobiol. Aging* 36, 3140–3151.
- Schneider, J.A., Arvanitakis, Z., Bang, W., Bennett, D.A., 2007. Mixed brain pathologies account for most dementia cases in community-dwelling older persons. *Neurology* 69, 2197–2204.
- Schneider, J.A., Arvanitakis, Z., Leurgans, S.E., Bennett, D.A., 2009. The neuropathology of probable Alzheimer disease and mild cognitive impairment. *Ann. Neurol.* 66, 200–208.
- Scott, J.A., Braskie, M.N., Tosun, D., Maillard, P., Thompson, P.M., Weiner, M., Decarli, C., Carmichael, O.T., 2016. Cerebral amyloid is associated with greater white-matter hyperintensity accrual in cognitively normal older adults. *Neurobiol. Aging* 48, 48–52.
- Shams, S., Granberg, T., Martola, J., Li, X., Shams, M., Fereshtehnejad, S.M., Cavallin, L., Aspelin, P., Kristoffersen-Wiberg, M., Wahlund, L.O., 2016. Cerebrospinal fluid profiles with increasing number of cerebral microbleeds in a continuum of cognitive impairment. *J. Cereb. Blood Flow Metab.* 36, 621–628.
- Skillback, T., Farahmand, B., Bartlett, J.W., Rosen, C., Mattsson, N., Nagga, K., Kilander, L., Religa, D., Wimo, A., Winblad, B., Rosengren, L., Schott, J.M., Blennow, K., Eriksdotter, M., Zetterberg, H., 2014. CSF neurofilament light differs in neurodegenerative diseases and predicts severity and survival. *Neurology* 83, 1945–1953.
- Smith, E.E., Greenberg, S.M., 2009. Beta-amyloid, blood vessels, and brain function. *Stroke J. Cereb. Circ.* 40, 2601–2606.
- Sudre, C.H., Cardoso, M.J., Bouvy, W.H., Biessels, G.J., Barnes, J., Ourselin, S., 2015. Bayesian model selection for pathological neuroimaging data applied to white matter lesion segmentation. *IEEE Trans. Med. Imaging* 34, 2079–2102.
- Tosto, G., Zimmerman, M.E., Hamilton, J.L., Carmichael, O.T., Brickman, A.M., 2015. The effect of white matter hyperintensities on neurodegeneration in mild cognitive impairment. *Alzheimer's Dement.* 11, 1510–1519.
- Van Westen, D., Lindqvist, D., Blennow, K., Minthon, L., N agga, K., Stomrud, E., Zetterberg, H., Hansson, O., 2016. Cerebral white matter lesions - associations with A $\beta$  isoforms and amyloid PET. *Scientific Rep.* 6, 1–9.
- Vemuri, P., Knopman, D.S., 2016. The role of cerebrovascular disease when there is concomitant Alzheimer disease. *Biochim. Biophys. Acta Mol. Basis Dis.* 1862, 952–956.
- Vemuri, P., Lesnick, T.G., Przybelski, S.A., Knopman, D.S., Lowe, V.J., Graff-Radford, J., Roberts, R.O., Mielke, M.M., Machulda, M.M., Petersen, R.C., Jack, C.R., 2017. Age, vascular health, and Alzheimer disease biomarkers in an elderly sample. *Ann. Neurol.* 82, 706–718.
- Vemuri, P., Lesnick, T.G., Przybelski, S.A., Knopman, D.S., Preboske, G.M., Kantarci, K., Raman, M.R., Machulda, M.M., Mielke, M.M., Lowe, V.J., Senjem, M.L., Gunter, J.L., Rocca, W.A., Roberts, R.O., Petersen, R.C., Jack, C.R., 2015. Vascular and amyloid pathologies are independent predictors of cognitive decline in normal elderly. *Brain* 138, 761–771.
- Verbeek, M.M., Kremer, B.P.H., Rikkert, M.O., Van Domburg, P.H.M.F., Skehan, M.E., Greenberg, S.M., 2009. Cerebrospinal fluid amyloid  $\beta$ 40 is decreased in cerebral amyloid angiopathy. *Ann. Neurol.* 66, 245–249.
- Wardlaw, J.M., Smith, E.E., Biessels, G.J., Cordonnier, C., Fazekas, F., Frayne, R., Lindley, R.I., O'Brien, J.T., Barkhof, F., Benavente, O.R., Black, S.E., Brayne, C., Breteler, M., Chabriat, H., Decarli, C., de Leeuw, F.E., Doubal, F., Duering, M., Fox, N.C., Greenberg, S., Hachinski, V., Kilimann, I., Mok, V., Oostenbrugge, R., Pantoni, L., Speck, O., Stephan, B.C., Teipel, S., Viswanathan, A., Werring, D., Chen, C., Smith, C., VAN Buchem, M., Norrving, B., Gorelick, P.B., Dichgans, M., Neuroimaging, S. T. F. R. V. C. O., 2013. Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration. *Lancet Neurol.* 12, 822–838.
- Weller, R.O., Subash, M., Preston, S.D., Mazanti, I., Carare, R.O., 2008. Perivascular drainage of amyloid- $\beta$  peptides from the brain and its failure in cerebral amyloid angiopathy and Alzheimer's disease. *Brain Pathol.* 18, 253–266.
- Zetterberg, H., Skillback, T., Mattsson, N., Trojanowski, J.Q., Portelius, E., Shaw, L.M., Weiner, M.W., Blennow, K., 2016. Association of cerebrospinal fluid neurofilament light Concentration with Alzheimer disease progression. *JAMA Neurol.* 73, 60–67.
- Zhou, B., Teramukai, S., Yoshimura, K., Fukushima, M., 2009. Validity of cerebrospinal fluid biomarkers as endpoints in early-phase clinical trials for Alzheimer's disease. *J. Alzheimers Dis.* 18, 89–102.
- Zhou, Y., Yu, F., Duong, T.Q., 2015. White matter lesion load is associated with resting state functional MRI activity and amyloid PET but not FDG in mild cognitive impairment and early Alzheimer's Disease patients. *J. Magn. Reson. Imaging* 41, 102–109.
- Zlokovic, B.V., 2011. Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nat. Rev. Neurosci.* 12, 723–738.