

HHS Public Access

Author manuscript *Hum Brain Mapp*. Author manuscript; available in PMC 2018 January 01.

Published in final edited form as:

Hum Brain Mapp. 2017 January ; 38(1): 151–164. doi:10.1002/hbm.23351.

Allometric scaling of brain regions to intra-cranial volume: an epidemiological MRI study

Laura W. de Jong, M.D^a, Jean-Sébastien Vidal, M.D. PhD^{b,C}, Lars E. Forsberg, MSc^d, Alex P. Zijdenbos, PhD^e, Thaddeus Haight, PhD^f, Alzheimer's Disease Neuroimaging Initiative^{*}, Sigurdur Sigurdsson, MSc^g, Vilmundur Gudnason, M.D. PhD^g, Mark A. van Buchem, M.D. PhD^a, and Lenore J. Launer, PhD^f

^aDepartment of Radiology, Leiden University Medical Center, Leiden, the Netherlands ^bAP-HP, Broca Hospital, Geriatrics department, Paris, 75013, France ^cUniversité Paris Descartes, Sorbonne Paris V, EA 4468, Paris, 75006, France ^dDepartment of Clinical Neuroscience Karolinska Institute, Stockholm, Sweden ^eBiospective Inc., Montreal, Canada ^fIntramural Research Program of the National Institute on Aging, Bethesda, Md USA ^gIcelandic Heart Association, Kopavogur, Iceland

Abstract

There is growing evidence that sub-structures of the brain scale allometrically to total brain size, i.e. in a non-proportional and non-linear way. Here, we examined scaling of different volumes of interest (VOI) to intra-cranial volume (ICV). It was assessed whether scaling was allometric or isometric and whether scaling coefficients significantly differed from each other. We also tested to what extent allometric scaling of VOI was introduced by the automated segmentation technique. Furthermore, reproducibility of allometric scaling was studied across different age groups and study populations. Study samples included samples of cognitively healthy adults from the community-based Age Gene/Environment Susceptibility-Reykjavik Study (AGES-Reykjavik Study) (N=3883), the Coronary Artery Risk Development in Young Adults Study (CARDIA) (N =709) and the Alzheimer's Disease Neuroimaging Initiative (ADNI) (N=180). Data encompassed participants with different age, ethnicity, risk factor profile, and ICV and VOI obtained with different automated MRI segmentation techniques. Our analysis showed that 1. Allometric scaling is a trait of all parts of the brain, 2. Scaling of neo-cortical white matter, neo-cortical gray matter, and deep gray matter structures including the cerebellum are significantly different from each other and 3. Allometric scaling of brain structures cannot solely be explained by age-associated atrophy, sex, ethnicity, or a systematic bias from study - specific segmentation algorithm, but appears to be a true feature of brain geometry.

Corresponding author L.W. de Jong, MD, Department of Radiology, Leiden University Medical Center, Albinusdreef 2, 2333 ZA Leiden, The Netherlands, Phone, + 31 71 529 9072, L.W.de_Jong@lumc.nl.

^{*}Some data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

The authors report no conflicts of interest.

Keywords

MRI; allometry; intra-cranial volume; brain; cortex; striatum; white matter; thalamus; AGES-Reykjavik; ADNI; CARDIA

1. Introduction

Since the development of (semi)-automated segmentation techniques for brain MRI, a large body of literature has emerged comparing brain volumes of different groups of people in order to find measurable traits distinctive or predictive for certain diseases. Having a good understanding of the physiologic variation in brain geometry is indispensable to discover pathological patterns. Human brain size varies considerably and different adjustment methods are applied to reduce noise stemming from this variation. Despite widespread use of standardization techniques, adjusting for ICV or total brain volume (TBV) when analyzing VOI is complex and controversial. In volumetric studies, ratios of VOI to ICV or TBV, or linear regression-based methods are commonly used. However, a critical evaluation of these techniques showed that each of these adjustment methods unmasks different types of relationships and results in different magnitude of effects (O'Brien, et al. 2011; Voevodskaya et al. 2014). In morphometric studies linear or non-linear stereotaxic registration of brain MR images are often used. A critical evaluation of these techniques showed that spatial transformation of MR brain images may result in significant opposite group level differences or different proportionality of brain regions compared to those obtained in native space (Allen, et al. 2008). Moreover, whether it is necessary to apply head-size adjustment in all types of comparative brain studies was evaluated in a study that investigated the effect of head size on several metrics of the brain, i.e. total brain volume, VOI, cortical thickness and voxel-based morphometry (VBM). It was concluded that head size adjustment should be considered in all volumetric and VBM studies, but not in cortical thickness studies (Barnes, et al. 2010).

Probably, part of the inconsistencies in results obtained with different head/ brain size adjustment methods can be explained by differences in underlying assumptions of these methods regarding preservation of proportionality of VOI to TBV across the total range of brain size variation in the population. Some techniques, such as ratio-based methods or linear registration, assume isometry of the brain, i.e. proportionality of VOI to TBV is preserved. Other techniques, such as linear regression-based methods or non-linear registration, allow for allometry to occur in case proportionality is not preserved. Although, these different theoretical underpinnings have been recognized (O'Brien, et al. 2011) and caution is called when choosing the adjustment method, it is uncertain whether allometrical scaling is true feature of brain geometry.

Some previous studies have provided evidence for allometric scaling of VOI to overall brain size. One study found larger proportions of cerebral WM and smaller proportions of GM in larger TBV compared to lower TBV (Luders, et al. 2002). Another study that focused on the necessity of head size, age and gender adjustment in MRI studies, found non linear relationships of cortical GM, hippocampus and putamen to ICV with a power less than 1

(Barnes, et al. 2010). Other neo-cortical metrics such as cortical thickness, total surface area, and sulcal depth have also been found to scale differently from what would be predicted based on ICV in case of isometry (Im, et al. 2008). Moreover, a recent study examined power law relationships of deep GM structures and many cortical GM regions and found most of them to have non-linear relationship with ICV. Some cortical areas had a power law larger than 1 and others smaller than 1. It was also tested whether prediction error of a statistical model would decrease when ICV correction was based on power-proportion method compared to the commonly used ANCOVA method. Prediction errors with use of power proportion method were slightly lower for structures that had strong non-linear relationships to ICV(Liu, et al. 2014).

Although, non-linearity and non-proportionality in scaling of some VOI to ICV have been reported, results are heterogeneous and little is known on scaling of especially deep GM regions (striatum and thalamus) and cerebellum. Also, it has not been investigated whether scaling coefficients of different brain structures are significantly different from each other. Here, scaling of volumes of frontal, parieto-occipital and temporal cortical GM, cortical WM, medial temporal lobe (MTL), striatum, thalamus, and cerebellum with ICV was studied using automatically segmented MRI brain scans of a large sample of community dwelling older adults (N = 3883) who participated in AGES-Reykjavik study. First, we investigated whether and to what extend VOI showed allometric scaling to ICV. Second, we estimated whether scaling coefficients of different VOI were significantly different from each other. Third, we studied whether scaling was similar in different age groups of our sample. Fourth, we set up an experiment to test whether the automated segmentation pipeline of AGES-Reykjavik Study could give rise to allometrical scaling. Fifth, because allometric scaling would have considerable influence on head/ brain size adjustment methods, the fit of the allometric model on the volumetric data was compared to the linear model. And lastly, since the AGES-Reykjavik study population consisted of older Icelandic individuals, extrapolation of our results to groups of younger individuals and/ or different ethnicity was potentially limited. Therefore, supportive analyses were conducted in two other samples (CARDIA and ADNI) that differed in mean age, source population, and method of automated MR segmentation to estimate brain volumes.

2. Materials and methods

2.1 General design of the AGES-Reykjavik Study

The general design and demographics of the AGES-Reykjavik have been described elsewhere (Harris, et al. 2007). The population-based sample of the AGES-Reykjavik consisted of 5764 men and women, born between 1907–1935. Participants underwent extensive clinical evaluation, including cognitive function testing and brain MRI. All participants signed an informed consent. The AGES-Reykjavik was approved by the Intramural Research Program of the National Institute on Aging, the National Bioethics Committee in Iceland (VSN00–063), the Icelandic Data Protection Authority, and the institutional review board of the U.S. National Institute on Aging, National Institutes of Health.

2.2 AGES-Reykjavik: MRI acquisition and automated MRI segmentation

MRI was performed at the Icelandic Heart Association on a single study dedicated 1.5-T GE Signa Twinspeed EXCITE system MRI scanner. The image protocol, described previously (Sigurdsson, et al. 2012), included a T1-weighted 3D spoiled gradient echo (TE 8 ms; TR 21 ms; FA 30°, FoV 240 mm; matrix 256×256 ; 110 slices; slice thickness 1.5 mm), a FSE PD/T2 (TE1 22 ms; TE2 90 ms; TR 3220 ms; echo train length 8; FA 90°, FoV 220 mm; matrix 256×256 ; slice thickness 3.0 mm), and a FLAIR (TE 100 ms; TR 8000 ms; inversion time 2000 ms; FA 90°, FoV 220 mm; matrix 256×256 ; slice thickness 3.0 mm).

A fully automated segmentation pipeline was developed based on the Montreal Neurological Institute processing pipeline (Sigurdsson, et al. 2012; Zijdenbos, et al. 2002). The pipeline used a multispectral approach to segment voxels into global tissue classes [cerebrospinal fluid (CSF), GM, WM and white matter hyperintensities (WMH)]. Following this, a regional parcellation pipeline - atlas-based segmentation method - was developed to obtain volumes of different sub-structures of the brain.

2.3 Determination of VOI

The regional tissue segmentation pipeline parcellated the brain in 56 different regions (Appendix A). However, for the present study, we combined regions into a limited amount of 8 VOI known to differ in gross cyto-architectural features. We separately assessed scaling of neo-cortical GM and WM to investigate in further detail the previously reported proportional changes as function of TBV. Three regions of neo-cortical GM were investigated, i.e. frontal (comprising of orbito-frontal and pre-frontal GM, precentral gyrus, cingulated gyrus, insula and fornix), temporal (comprising of lateral temporal GM, parahippocampal and fusiform gyrus), and parieto-occipital GM. Cortical WM volume was studied in total and included all lobar WM, corpus callosum, internal and external capsule, and WMH. The medial temporal lobe (MTL), striatum, thalamus and cerebellum were separately studied because of their importance in many studies to neurodegenerative processes. MTL included amygdala and hippocampus (including CA regions I-IV, fimbria, and subiculum of the hippocampus). Striatum included the nucleus accumbens, caudate nucleus, putamen, and globus pallidus. The thalamus included also the hypothalamus. The cerebellum included cerebellar GM and WM. Left and right hemispheres of each structure were combined. Total brain volume (TBV) was calculated as the sum of the neo-cortical GM and WM, MTL, striatum, thalamus, brainstem and cerebellum. ICV was defined as the sum of TBV and CSF.

2.4 Quality control of tissue classification and validation of VOI

The quality of the segmentation of the 8 composite VOI was mostly dependent on the performance of the global tissue segmentation into GM, WM, WMH, and CSF, and for a small part dependent on the definition of topographical borders by the regional parcellation pipeline. Performance of both global tissue segmentation and regional tissue parcellation was evaluated. The quality control of global tissue classification consisted of 3 steps described in (Sigurdsson, et al. 2012). In summary these were: 1) Visual inspection of the segmentation of 14 a priori selected slices of each subject (N= 4356), which led to additional manual editing in 43 cases and rejection of 53 cases. 2) Comparison of automated versus

manual global tissue segmentation of 5 preselected slices across the brain (including a slice located at the junction of the thalamus and subthalamic structures for reviewing segmentation of the deep gray matter nuclei) in 20 randomly selected cases. Resulting dice similarity index scores (Zijdenbos, et al. 1994) were 0.82, 0.82, and 0.83 for GM, WM, and CSF respectively. 3) Reproducibility of the entire process of MRI acquisition and postprocessing was evaluated by repeated scanning and segmentation (4 times in total) of 32 participants. Excellent intra-class correlation for all global tissue was found (r > 0.98, for all). Because the present study relies for an important part on good quality of ICV segmentation, the performance of the automated pipeline was further evaluated specifically on ICV. ICV was manually segmented on the same 20 brain scans used for step 2 of the quality control. Two researchers with extensive neuroradiological experience and blinded for the results of the automated segmentation, segmented ICV on axial 3D T1 weighted images, with correction and editing in sagittal and coronal planes. Resulting ICV were correlated to ICV obtained by the automated pipeline. Pearson's correlation was 0.97 (0.93–0.99) and Bland-Altmann plot showed a small overestimation of ICV of 31 cm³ on average by the automated segmentation, but no proportional error (Appendix B and C).

Performance of regional parcellation pipeline was validated against four complete manually labeled scans. Dice similarity index scores per studied region were; frontal GM: 0.83, temporal GM: 0.83, parieto-occipital GM: 0.81, striatum: 0.83, MTL: 0.80, thalamus: 0.92, cerebellum: 0.92, white matter: 0.86.

2.5 Statistical analysis

All statistical analyses were performed with SAS v 9.13 (SAS Institute Inc., Cary, NC, USA) and all graphs were generated with R v 3.1.2 (R Core Team (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/).

2.5.1 Analytical sample of AGES - Reykjavik Study—MR scanning was performed on consenting MR eligible participants, between 2002 and 2006. From the total AGES-Reykjavik sample of 5764 participants, 4726 underwent successful MRI scanning. Global and regional segmentations were successful in 4613 MR scans. We excluded cases of dementia (N = 202) and MCI (N = 422), assumed to have higher rates of atrophy, and cases for which cognitive function had not been assessed (N=106). Our final study sample consisted of 3883 people with successful brain MRI and segmentation of the images. Demographics and brain structure volumes of the AGES-Reykjavik study population were compared between women and men with t-tests for continuous variables and chi-square tests for categorical variables. All VOI were normally distributed.

2.5.2 Estimation of scaling coefficients of different VOI—Allometric coefficients of VOI with ICV were calculated using the general equation for allometric analyses, log(y) = log(b) + a log(x), where x is ICV, y VOI, log(b) intercept, and a represents the allometric coefficient (Harvey 1982), i.e. the slope of the regression between log (ICV) and log (VOI). A coefficient greater than 1.0 is considered a positive allometric coefficient, i.e. VOI increased with a power greater than 1 relative to ICV. A coefficient smaller than 1.0 is seen

as a negative allometric coefficient, i.e. VOI increased with a power less than 1 relative to ICV.

We chose ICV, instead of TBV, as measure of brain volume to avoid a possible bias towards isometry in estimating allometric coefficients of large VOI. Large structures occupy large volumes in TBV making the range of possible deviations from isometry smaller; this may produce an overestimate of coefficients towards 1 and reduce the ability to estimate allometric coefficients deviant from 1 (Deacon 1990). With the use of ICV none of the structures studied comprised more than 24% (WM) of ICV. Another important reason was that ICV is regarded as a marker for brain volume at its maximum size and therefore a marker of "pre-morbid" brain size. At time of scanning, brains of most study participants experienced more or less atrophy due to aging or pathological processes. These are factors we can largely control for in our statistical analyses, whereas it is more difficult to control for differences between current TBV without taking into account the original size of the brain at maximum. Log-transformed VOI were plotted against log transformed ICV (figure 1). For each VOI, allometric coefficients with ICV were calculated adjusted for age and sex, (log (VOI) = intercept + α log (ICV) + $\beta_{age} \times age + \beta_{sex} \times sex$) and tested against the isometric scaling law of 1:1.

2.5.3 Comparison of allometric scaling coefficients of different VOI—Allometric coefficients of the different VOI to ICV were compared using a marginal model (PROC MIXED SAS procedure (SAS Institute Inc., Cary, NC, USA) with repeated statement and unstructured correlation matrix), which takes into account the correlations between the VOI. The log transformed VOI were entered as dependent variables and log transformed ICV as independent variable. Interactions of log(VOI) with log(ICV) were entered in the model as a cross product together with log(ICV), log(VOI), age, and sex. The model was also run with additional independent variables (year of birth, height, achievement of higher education (highschool diploma or above), presence of infarct(s) yes/no, and contrast-to-noise ratio (CNR) between GM and WM and CNR between GM and cerebro-spinal fluid), but these did not exert significant effects and were omitted to keep the model parsimonious. A Bonferroni correction was applied to adjust for multiple testing (number of comparisons between slopes in the 3 mixed models = 85) and a p-value < 0.00059 (=0.05/85) was considered significant. The analysis was performed in the entire sample and repeated for women and men separately. The numerical results of the marginal model are reported in table 2.

2.5.4 Allometric scaling of VOI in different age groups—To assess whether age influenced scaling of VOI with ICV, scaling coefficients of VOI to ICV were calculated for each quartile of age; the age range of the youngest quartile being 66–71 years, and of subsequent quartiles being, 72–75, 76–79 and 80–95 years. The coefficients were compared among the quartiles by testing whether there was an interaction between the quartiles and ICV.

2.5.5 Testing the automated segmentation pipeline with artificially linearly scaled data—To test whether a potential systematic error in the automated segmentation pipeline could introduce allometry in the volumetric data of AGES-RS, artificially linearly scaled brain scans were entered into the pipeline and the output was investigated for

allometry. Scans of a relatively small (1402 cm³) and relatively large brain (1756 cm³) were skull stripped and linearly scaled by factors ranging from 0.75–1.25 of its original size with steps of 0.01. The resulting sets of scaled images were subsequently processed through the AGES-Reykjavik pipeline. Log transformed volumes of the global tissues GM, WM and CSF were plotted against log transformed ICV and α -coefficients were calculated.

2.5.6 Comparison of allometric model and linear regression model—The fit of the allometric model of the relationship of each VOI to ICV on the data was compared to a linear regression model. The line of prediction from the allometric model and linear model were superimposed in the same graph and R^2 of each model was calculated. Both models were conducted with adjustments for age and sex.

2.5.7 Allometric scaling in different study populations; supportive analyses in datasets of CARDIA and ADNI—Supportive analyses were conducted in datasets of CARDIA and ADNI. In both samples the allometric coefficients of VOI with ICV were calculated, corrected for age and sex, and tested against the isometric scaling law of 1:1, similar to the first part of analysis conducted in the AGES-Reykjavik data.

The multi-center prospective cohort CARDIA study was designed to examine the development and determinants of clinical and subclinical cardiovascular disease and its risk factors. Between 1985–1986, 5115 black and white men and women (aged 16–30) were recruited from 4 urban sites across the United States and underwent 8 examination cycles (Friedman, et al. 1988). All participants provided written informed consent at each exam, and institutional review boards from each study site and the coordinating center annually approved the study. In 2010-2011, 3498 (72%) of the surviving cohort attended a 25-year follow-up exam. As part of this exam, a sub-sample of the cohort participated in the CARDIA Brain sub-study, designed to investigate the morphology, pathology, physiology and function of the brain with MRI. Exclusion criteria at the time of sample selection, or at the MRI site, were a contra-indication to MRI or a body size that was too large for the MRI scanner. Of those who were eligible for the sub-study, 719 individuals received whole brain MRI scans. Post-scan image processing was performed by the Section of Biomedical Image Analysis (BIA), Department of Radiology, University of Pennsylvania. MRI scans were inspected and passed through a quality control process. Based on previously described methods (Davatzikos, et al. 2003; Goldszal, et al. 1998; Lao, et al. 2008; Shen, et al. 2002; Zacharaki, et al. 2008), an automated algorithm was used to segment MRI structural images of supratentorial brain tissue into GM, WM and cerebrospinal fluid. GM and WM were further characterized and segmented as 92 anatomic ROIs in each hemisphere, from which summary VOIs used in the current study were calculated. ICV was calculated as the sum of all supratentorial structures, but not infratentorial.

Some data used in the preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in 2003 by the National Institute on Aging (NIA), the National Institute of Biomedical Imaging and Bioengineering (NIBIB), the Food and Drug Administration (FDA), private pharmaceutical companies and non-profit organizations, as a \$60 million, 5-year public-private partnership. The primary goal of ADNI has been to test whether serial

magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). Determination of sensitive and specific markers of very early AD progression is intended to aid researchers and clinicians to develop new treatments and monitor their effectiveness, as well as lessen the time and cost of clinical trials.

The Principal Investigator of this initiative is Michael W. Weiner, MD, VA Medical Center and University of California – San Francisco. ADNI is the result of efforts of many coinvestigators from a broad range of academic institutions and private corporations, and subjects have been recruited from over 50 sites across the U.S. and Canada. The initial goal of ADNI was to recruit 800 subjects but ADNI has been followed by ADNI-GO and ADNI-2. To date these three protocols have recruited over 1500 adults, ages 55 to 90, to participate in the research, consisting of cognitively normal older individuals, people with early or late MCI, and people with early AD. The follow up duration of each group is specified in the protocols for ADNI-1, ADNI-2 and ADNI-GO. Subjects originally recruited for ADNI-1 and ADNI-GO had the option to be followed in ADNI-2. For up-to-date information, see www.adni-info.org.

For the supportive analysis of our study we used volumetric brain measures derived from the standardized 1.5 T MRI screening dataset in cognitively healthy subjects that was collected between August 2005 and October 2007 and processed using FreeSurfer software (Freesurfer Software Site. Cortical Reconstruction and volumetric segmentation, (http://surfer.nmr.mgh.harvard.edu/; 2016 [accessed 07.26.16]).

3. Results

3.1 Characteristics of AGES-Reykjavik sample

The AGES-Reykjavik sample had a mean age of 75.7 (standard deviation = 5.2) years, of which 59.8% were women. ICV ranged from 1116 cm³ – 2162 cm³ in the total sample; 1116 cm³ – 1868 cm³, in women and 1232 cm³ – 2162 cm³ in men. Women had on average lower educational level (p < 0.0001), higher BMI (p = 0.003), were diagnosed less often with diabetes (p < 0.0001), and had smoked (p < 0.0001) and drank alcohol (p < 0.0001) more sparingly compared to men. Women had lower means of ICV and all VOI compared to men (p < 0.0001) (table 1).

3.2 Allometric scaling coefficients of all VOI

All VOI scaled non-isometrically to ICV (figure 1). After correction for age and sex, a positive allometric coefficient of 1.14 (95% confident interval=1.11–1.17)) was estimated for WM volume and negative allometric coefficients were found for frontal GM [0.76 (0.73–0.79)], temporal GM [0.75 (0.72–0.78)], parieto-occipital GM [0.79 (0.76–0.83)], MTL [0.60 (0.56–0.64)], thalamus [0.59 (0.56–0.62)], striatum [0.41 (0.37–0.45)], and cerebellum [0.55 (0.52–0.59)]. All were found significantly differently from 1 (1:1 scaling law to ICV (p < 0.0001)).

3.3 Significant scaling differences between VOI

Results from the marginal model showed that the α -coefficient of WM volume to ICV was significantly different from the α -coefficients of all GM VOI (table 2) in the entire sample, and in women and men separately. The α -coefficients of the different neo-cortical GM areas to ICV were not significantly different from each other in women and men separately. Also, the α -coefficients of MTL, thalamus, and cerebellum were not significantly different from each other in women and men separately. However, in the entire sample the α -coefficient of the MTL was not significantly different from the α -coefficient of the parieto-occipital GM, but was significantly different from the thalamus and the cerebellum. The α -coefficient of the striatum was significantly different from all other α -coefficients except for the α -coefficient of the thalamus and cerebellum in the entire sample, and the α -coefficient of the cerebellum in men only.

3.4 Allometric scaling in different age groups

a-coefficients of VOI to ICV for each quartile of age are shown in table 3. a-coefficients of the both cortical and deep GM structures and cerebellum in the older quartiles appeared somewhat lower compared to the younger quartiles and the a-coefficient of WM appeared higher in the older quartiles. However, these differences were non-significant, except for temporal GM which was significantly lower in the older quartiles compared to the younger (*p*-value of 0.004).

3.5 Little allometry introduced by automated segmentation pipeline

Figure 2 displays log of global tissue volumes plotted against log ICV obtained by the automated segmentation pipeline based on the artificially linearly scaled data set of a relatively large and small brain. The α -coefficients were 0.99 for GM, 1.02 for WM, and 1.00 for CSF for dataset based on the relatively large brain and 0.98 for GM, 1.01 for WM, and 10.2 for CSF for the dataset based on the relatively small brain. Because of the almost perfect fit of the points and the regression line these α -coefficients were significantly different from the isometric scaling law of 1.0 (all *p*-values < 0.0001), except for the CSF in the large brain.

3.6 Comparable fit of allometric model and linear regression model

Figure 3 superimposes the line of prediction of the allometric model (and associated α coefficient and R^2) with line of prediction of the linear model (and associated β -coefficient and R^2). Compared to the R^2 of the linear model, the R^2 of the allometric model was a few per mille smaller for cerebellar, cortical and deep GM structures and a few per mile larger for WM. The models have a comparable fit and can substitute each other.

3.7 Allometric scaling in CARDIA and ADNI

The CARDIA sample consisted of individuals with a mean age of 50 (3.5) years, of which 52.9% were women. ICV in the CARDIA sample, including only supratentorial areas, varied from 999 cm³ – 1643 cm³. The ADNI sample consisted of individuals with a mean age of 76 (5.0) years, of which 49.4% were women. ICV in the ADNI sample varied from 1116 cm³ – 1985 cm³. We found the highest allometric coefficients for WM volume in both CARDIA (α

= 1.05) and ADNI (α = 1.00). All GM areas had negative allometric coefficients with the lowest coefficients in the deep GM areas (table 3). Roughly, results suggest similar trends compared with those found in the AGES Reykjavik data. However, important differences were 1) allometric coefficients of the neo-cortical GM areas to ICV in CARDIA with values between 0.90–0.94 were higher, compared with those in the AGES and ADNI samples, and 2) WM volume in the ADNI data set seems to increase isometrically with ICV.

4. Discussion

Different allometric scaling of neo-cortical WM, neo-cortical GM and deep GM

One goal of the present study was to assess and compare scaling coefficients of different VOIs to ICV in the AGES-Reykjavik dataset. We found all VOI to scale allometrically with ICV. One could roughly discern three patterns of scaling, i.e. WM scaling, neocortical GM scaling and deep GM scaling. First, neo-cortical WM was the only structure to proportionally increase in larger ICV with a positive allometric coefficient of 1.14. Scaling of WM was found significantly different from all GM structures and cerebellum. Second, negative allometric coefficients were found for frontal (0.76), temporal (0.74), and parieto-occipital (0.79) cortical GM structures. Scaling coefficients of neo-cortical GM structures (frontal, temporal, and parieto-occipital) were not significantly different from each other, but were significantly larger than scaling coefficients of the deep GM structures when women and men were separately assessed. Also scaling coefficients of MTL (0.60), thalamus (0.59), and cerebellum (0.55) were not significantly different from each other in women and men separately.

Allometric scaling cannot solely be explained by age, sex, ethnicity or a systemic bias from segmentation pipeline

One important limitation of our study was that the sample consisted of older individuals, who have experienced various amounts of brain atrophy. Therefore, the observed scaling coefficients cannot be extrapolated to younger samples. After stratifying the AGES Reykjavik sample into quartiles of age, we found most structures to have similar scaling coefficients except for temporal GM (not including the MTL), which had lower α -coefficients in older individuals. We do not have an explanation for the significant difference in scaling found for temporal GM, but it prompted us not to rule out the possibility that allometric scaling of sub structures of the brain may vary with age in a way that we could not detect in the age span of our sample. Nonetheless, the findings of this paper show that allometric scaling is a feature of the brain in the older population, which cannot be accounted for by adjusting for age when performing brain comparative studies.

A second important limitation was that all participants were Icelandic and the sample was genetically relatively homogeneous. Ancillary analyses in ADNI and CARDIA, with participants of younger age and different ethnicities, also showed that WM increased proportionally steepest with increasing ICV, followed by proportionally decreases in GM, with greatest decreases in the deep GM structures, similar to our observations in the AGES Reykjavik Study. Still, there were also differences in the results between the studies. The allometric coefficients of the cortical GM areas to ICV in CARDIA seemed higher

compared those in the AGES and ADNI samples. A potential explanation could be that "ICV" in CARDIA was constructed from supratentorial structures only, and as a result allometric coefficients were higher. Another explanation could be that in the relatively young sample of the CARDIA allometric scaling is less pronounced. Further studies are needed to specifically examine this hypothesis. A second difference between the results of the additional analyses and our primary analyses in AGES was that WM volume in the ADNI data set seemed to increase isometrically with ICV. This may be explained by differences in tissue segmentation between GM and WM, as suggested by the higher mean volume of WM and lower mean volume of GM in ADNI compared with AGES Reykjavik. Depending on how border voxels are assigned to the GM and WM tissue classes, the difference between allometric coefficients may differ. Because of these differences in scaling coefficients among the study samples, it is at the moment not possible to establish fixed reproducible allometric coefficients for the human brain and more studies are needed.

A third potential limitation of the study was the use of an automated MR segmentation technique. Systematic errors, such as improper skull stripping, incorrect intensity thresholds, difficulty in segmenting sulcal CSF or imprecise template warping could all be possible sources of finding allometric correlations between VOI to ICV. However, when we fed artificially linearly scaled scans in the segmentation pipeline, the scaling coefficients of the output only showed small deviations from the isometric scaling law of 1, at maximum in the order of 2 %. This could not explain the much larger deviations from 1 of the different scaling laws of VOI in the study sample. Therefore, we did not find evidence for a possible systematic error in the segmentation pipeline that could explain the allometry.

Allometric scaling as true feature of brain geometry

Differences in geometric or cyto-architectural properties of different brain structures may underlie differences and similarities in scaling to ICV. We observed similar scaling coefficients of different neo-cortical GM areas, which suggest they preserve proportionality to one another regardless of ICV. However, cortical GM and WM had significantly different scaling coefficients, indicating they do not preserve proportionality with varying brain size. This can be explained by differences in topology, where GM can be regarded as a surface of neural tissue covering an associated volume of WM (Dale, et al. 1999). The different lobes of the neo-cortex are similarly organized in repetitive cortical columns (Mountcastle 1997). Assuming a stable thickness of the neo-cortical GM "surface" across various brain sizes, as suggested by several studies (Hofman 1985; Hofman 1988; Mountcastle 1997), neo-cortical GM to WM should scale by an exponent of 2/3 (square-cube). If we focus on the results based on the AGES-Reykjavik study sample, we can observe that scaling coefficients of neocortical white to gray matter range from 0.65 to 0.70 (0.76/1.14 for frontal GM, 0.74/1.14 for temporal GM, and 0.79/1.14 for parieto-occipital GM), which approximates the geometric square-cube scaling law. Nevertheless, we did not establish the same results in the younger sample of CARDIA or the smaller sample of the ADNI and caution should be taken to apply a purely square-cube scaling law to the architecture of neo cortex. In a previous study slight increases of neo-cortical thickness (scaling of 0.2) with increase in ICV were observed (Im, et al. 2008). Another recent study showed the neo-cortical GM to have a more extensive gyrification, i.e. to be "twistier", in larger brains compared to smaller ones

(Germanaud, et al. 2012). Also, for other parameters, such as cell soma size or amount of supporting glial cells, the extent to which they vary with increasing brain size is unclear. Some studies have also pointed to possible constraints in WM expansion, which should lead to scaling factors of white to gray that are higher than the square-cube law of 2/3. It has been proposed that hemispheric specialization increases with increasing brain volume, which would lead to a decrease in inter-hemispheric connections and thus a decrease in WM volume (Ringo, et al. 1994). However, the coefficients reported in the present study for the AGES-Reykjavik study provide no evidence for such a limitation on WM expansion

The disproportionally lower scaling coefficients of deep GM structures to ICV compared to the cortex are not readily explained. The cortex gives rise to connections with striatum and thalamus, thus these structures could be expected to expand with neo-cortical GM volume. However, we found no evidence for preserved proportionality of the striatum and thalamus with cortical GM with scaling. Possibly, the deep GM structures are also influenced by other factors during brain development than neo-cortical growth. Brain structures grow in asynchronous patterns from birth through early adulthood and the striatum has been shown, together with frontal brain areas, to undergo more extensive developmental changes relatively late in early adulthood compared to other brain areas (Sowell, et al. 1999). Also, genetic factors could influence variation in regional brain volumes and lead to disproportional neo-cortical and deep GM volume increases with ICV, especially for the striatum. One twin study showed that the volume of the striatum, thalamus, and cerebellum were significantly more influenced by genetic factors compared to neo-cortical structures that were influenced more by environmental factors (Yoon, et al.). And another twin-study concluded the phenotypic covariance of the striatal structures, hippocampus, and thalamus was primarily due to patterns of genetic covariance (Eyler, et al.).

Implications of allometric scaling for methods of head/ brain size adjustment

Knowledge on the allometric scaling of regional brain volumes is important for the discussion of adjustment methods for normal variation in comparative brain studies to volumetric and morphological changes. Allometric scaling implies both non-proportionality and non-linearity of scaling. Our results contribute to the understanding why certain methods should not be used. Ratios of brain structure volume over ICV or stereotaxic normalization by means of linear affine transformation assume isometric scaling of the brain, i.e. proportionate scaling, which may lead to over- or underestimation of results. Therefore the use of these methods should be avoided, except when studying disproportionality is the purpose. The erroneous effect of linear spatial normalization on groups with differences in ICV was illustrated in a study comparing neo-cortical thickness differences in stereotaxic and native space between men and women (Luders, et al. 2006). The normalized data showed a disproportionately increased neo-cortical thickness in women compared to men, which was considerably attenuated in the unscaled data. Another important finding of our study was that allometric scaling was most apparent in deep GM structures. Unwanted effects of spatial registration therefore may be expected to be especially problematic in deep gray matter structures. Previously, a study reported that spatial-transformation based methods indeed produce significantly different proportions in smaller structures such as the hippocampus (Allen, et al. 2008). Lastly, we compared the fit

of the allometric model to a linear model in predicting the relationship of VOI to ICV. We found very small differences in \mathbb{R}^2 , which implies the allometric model and linear model could substitute each other in the range of total brain size variation among humans. Therefore, we conclude that it is important in brain comparative studies to adjust for non-proportionality, but not for non-linearity.

5. Conclusion

In summary, our study found allometric scaling of WM, neo-cortical GM and deep GM structures to ICV in large samples of adult humans with different age, sex and ethnicity. A positive allometric coefficient was found for WM and negative allometric coefficients for neo-cortical and deep GM structures, with smallest scaling coefficients for deep GM. Furthermore, our analysis showed that the allometric scaling cannot solely be explained by age, sex, ethnicity, or a possible systematic bias arising from the automated segmentation algorithm. We therefore conclude allometry is a true feature of the brain geometry.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to thank the participants of the AGES-Reykjavik Study and the IHA clinic staff for their invaluable contribution.

Funding

This study was funded by the National Institute on Aging Intramural Research Program (N01-AG-12100), Hjartavernd (the Icelandic Heart Association), and the Althingi (the Icelandic Parliament). The study was approved by the Icelandic National Bioethics Committee (VSN: 00-063) and the Medstar Research Institute (project 2003-145). Dr. de Jong is supported by the Netherlands Organisation for Health Research and Development (AGIKO grant number 92003536). Prof van Buchem is supported by an unrestricted grant from the Dutch Genomics Initiative (NCHA 050-060-810). Drs. Vidal, Zijdenbos, Gudnason and Launer, and Mr. Sigursson report no disclosures.

ADNI

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen Idec Inc.; Bristol-Myers Squibb Company; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; ; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Medpace, Inc.; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Synarc Inc.; and Takeda Pharmaceutical Company. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of Southern California.

CARDIA

The Coronary Artery Risk Development in Young Adults Study (CARDIA) is supported by contracts HHSN268201300025C, HHSN268201300026C, HHSN268201300027C, HHSN268201300028C,

HHSN268201300029C, and HHSN268200900041C from the National Heart, Lung, and Blood Institute (NHLBI), and an intra-agency agreement between NIA and NHLBI (AG0005).

Abbreviations

ICV	intra-cranial volume
TBV	total brain volume
MRI	magnetic resonance imaging
AGES-Reykjavik	Age Gene/Environment Susceptibility-Reykjavik Study
GM	gray matter
WM	white matter
MTL	medial temporal lobe
CARDIA	The Coronary Artery Risk Development in Young Adults Study
ADNI	Alzheimer's Disease Neuroimaging Initiative

References

Allen JS, Bruss J, Mehta S, Grabowski T, Brown CK, Damasio H. Effects of spatial transformation on regional brain volume estimates. Neuroimage. 2008; 42(2):535–547. [PubMed: 18599317]

- Barnes J, Ridgway GR, Bartlett J, Henley SM, Lehmann M, Hobbs N, Clarkson MJ, MacManus DG, Ourselin S, Fox NC. Head size, age and gender adjustment in MRI studies: a necessary nuisance? Neuroimage. 2010; 53(4):1244–1255. [PubMed: 20600995]
- Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. Neuroimage. 1999; 9(2):179–194. [PubMed: 9931268]
- Davatzikos C, Tao X, Shen D. Hierarchical active shape models, using the wavelet transform. IEEE Trans Med Imaging. 2003; 22(3):414–423. [PubMed: 12760558]
- Deacon T. Fallacies of progression in theories of brain-size evolution. International Journal of Primatology. 1990; 11(3):193–236.
- Eyler LT, Prom-Wormley E, Fennema-Notestine C, Panizzon MS, Neale MC, Jernigan TL, Fischl B, Franz CE, Lyons MJ, Stevens A, Pacheco J, Perry ME, Schmitt JE, Spitzer NC, Seidman LJ, Thermenos HW, Tsuang MT, Dale AM, Kremen WS. Genetic patterns of correlation among subcortical volumes in humans: Results from a magnetic resonance imaging twin study. Hum Brain Mapp. 2010; 32(4):641–653.
- Friedman GD, Cutter GR, Donahue RP, Hughes GH, Hulley SB, Jacobs DR Jr, Liu K, Savage PJ. CARDIA: study design, recruitment, and some characteristics of the examined subjects. J Clin Epidemiol. 1988; 41(11):1105–1116. [PubMed: 3204420]
- Germanaud D, Lefevre J, Toro R, Fischer C, Dubois J, Hertz-Pannier L, Mangin JF. Larger is twistier: spectral analysis of gyrification (SPANGY) applied to adult brain size polymorphism. Neuroimage. 2012; 63(3):1257–1272. [PubMed: 22877579]
- Goldszal AF, Davatzikos C, Pham DL, Yan MX, Bryan RN, Resnick SM. An image-processing system for qualitative and quantitative volumetric analysis of brain images. J Comput Assist Tomogr. 1998; 22(5):827–837. [PubMed: 9754125]
- Harris TB, Launer LJ, Eiriksdottir G, Kjartansson O, Jonsson PV, Sigurdsson G, Thorgeirsson G, Aspelund T, Garcia ME, Cotch MF, Hoffman HJ, Gudnason V. Age, Gene/Environment Susceptibility-Reykjavik Study: multidisciplinary applied phenomics. Am J Epidemiol. 2007; 165(9):1076–1087. [PubMed: 17351290]

Harvey PH. On rethinking allometry. J Theor Biol. 1982; 95(1):37-41. [PubMed: 7087496]

- Hofman MA. Size and shape of the cerebral cortex in mammals. I. The cortical surface. Brain Behav Evol. 1985; 27(1):28–40. [PubMed: 3836731]
- Hofman MA. Size and shape of the cerebral cortex in mammals. II. The cortical volume. Brain Behav Evol. 1988; 32(1):17–26. [PubMed: 3056571]
- Im K, Lee JM, Lyttelton O, Kim SH, Evans AC, Kim SI. Brain size and cortical structure in the adult human brain. Cereb Cortex. 2008; 18(9):2181–2191. [PubMed: 18234686]
- Lao Z, Shen D, Liu D, Jawad AF, Melhem ER, Launer LJ, Bryan RN, Davatzikos C. Computerassisted segmentation of white matter lesions in 3D MR images using support vector machine. Acad Radiol. 2008; 15(3):300–313. [PubMed: 18280928]
- Liu D, Johnson HJ, Long JD, Magnotta VA, Paulsen JS. The power-proportion method for intracranial volume correction in volumetric imaging analysis. Front Neurosci. 2014; 8:356. [PubMed: 25414635]
- Luders E, Narr KL, Thompson PM, Rex DE, Woods RP, Deluca H, Jancke L, Toga AW. Gender effects on cortical thickness and the influence of scaling. Hum Brain Mapp. 2006; 27(4):314–324. [PubMed: 16124013]
- Luders E, Steinmetz H, Jancke L. Brain size and grey matter volume in the healthy human brain. Neuroreport. 2002; 13(17):2371–2374. [PubMed: 12488829]
- Mountcastle VB. The columnar organization of the neocortex. Brain. 1997; 120(Pt 4):701–722. [PubMed: 9153131]
- O'Brien LM, Ziegler DA, Deutsch CK, Frazier JA, Herbert MR, Locascio JJ. Statistical adjustments for brain size in volumetric neuroimaging studies: some practical implications in methods. Psychiatry Res. 2011; 193(2):113–122. [PubMed: 21684724]
- Ringo JL, Doty RW, Demeter S, Simard PY. Time is of the essence: a conjecture that hemispheric specialization arises from interhemispheric conduction delay. Cereb Cortex. 1994; 4(4):331–343. [PubMed: 7950307]
- Shen D, Moffat S, Resnick SM, Davatzikos C. Measuring size and shape of the hippocampus in MR images using a deformable shape model. Neuroimage. 2002; 15(2):422–434. [PubMed: 11798276]
- Sigurdsson S, Aspelund T, Forsberg L, Fredriksson J, Kjartansson O, Oskarsdottir B, Jonsson PV, Eiriksdottir G, Harris TB, Zijdenbos A, van Buchem MA, Launer LJ, Gudnason V. Brain tissue volumes in the general population of the elderly: the AGES-Reykjavik study. Neuroimage. 2012; 59(4):3862–3870. [PubMed: 22119006]
- Sowell ER, Thompson PM, Holmes CJ, Jernigan TL, Toga AW. In vivo evidence for post-adolescent brain maturation in frontal and striatal regions. Nat Neurosci. 1999; 2(10):859–861. [PubMed: 10491602]
- Voevodskaya O, Simmons A, Nordenskjold R, Kullberg J, Ahlstrom H, Lind L, Wahlund LO, Larsson EM, Westman E. The effects of intracranial volume adjustment approaches on multiple regional MRI volumes in healthy aging and Alzheimer's disease. Front Aging Neurosci. 2014; 6:264. [PubMed: 25339897]
- Yoon U, Perusse D, Lee JM, Evans AC. Genetic and environmental influences on structural variability of the brain in pediatric twin: deformation based morphometry. Neurosci Lett. 2011; 493(1–2):8– 13. [PubMed: 21296128]
- Zacharaki EI, Kanterakis S, Bryan RN, Davatzikos C. Measuring brain lesion progression with a supervised tissue classification system. Med Image Comput Comput Assist Interv. 2008; 11(Pt 1): 620–627. [PubMed: 18979798]
- Zijdenbos AP, Dawant BM, Margolin RA. Automatic detection of intracranial contours in MR images. Comput Med Imaging Graph. 1994; 18(1):11–23. [PubMed: 8156533]
- Zijdenbos AP, Forghani R, Evans AC. Automatic "pipeline" analysis of 3-D MRI data for clinical trials: application to multiple sclerosis. IEEE Trans Med Imaging. 2002; 21(10):1280–1291. [PubMed: 12585710]

Appendices

Appendix A, Brain regions of the probabilistic atlas

Regions of interest:

- Striatum:1 caudate nucleus, 2 putamen, 3 globus pallidus
- Thalamus: 4
- Medial temporal lobe: 5 amygdala, 6 hippocampus
- Frontal GM: 7 cingulate gyrus, 8 prefrontal lobe, 9 precentral gyrus, 10 insula
- Parieto-occipital GM: 12 occipital lobe, 14 parietal lobe
- Temporal GM: 11 lateral temporal lobe, 13 parahippocampal gyrus, 15 fusiform gyrus
- Cerebellum: 30

Appendix B, Accuracy of automated segmentation pipeline; Pearson's correlation manual versus automated segmentation of ICV

Appendix C, Accuracy of automated segmentation pipeline; Bland-Altman plot manual versus automated segmentation of ICV

Appendix D, Dice similarity index scores for regional tissue classification

de Jong et al.

Page 17



Figure 1.

Allometric coefficients of VOI with ICV

Gray line, isometry line; Red line, line of the allometric log-log model between the ICV and the VOI.

de Jong et al.



Figure 2. Accuracy of automated segmentation pipeline; scaling of artificially linearly scaled data

de Jong et al.

Page 19



Figure 3.

Comparison of allometric log-log model to linear model of VOI to ICV Red line, line of the allometric log-log model between the ICV and the VOI; Blue line, line of the linear model between the ICV and the VOI.

Table 1

General characteristics of the study sample

Mean (SD)*	All (N=3883)	Women (N=2307)	Men (N=1576)	\mathbf{p}^{\dagger}
Age [years]	75.7 (5.2)	75.6 (5.3)	75.8 (5.1)	0.27
Higher education, % (N)	12.2 (473)	6.52 (150)	20.6 (323)	< 0.0001
BMI [kg/m ²]	27.0 (4.3)	27.2 (4.7)	26.8 (3.7)	0.003
Diabetes, % (N)	11.1 (430)	8.76 (202)	14.5 (228)	< 0.0001
Smoking status, % (N)				
Never	41.7 (1619)	53.2 (1226)	24.9 (393)	
Former	44.5 (1728)	34.8 (803)	58.7 (925)	< 0.0001
Current	13.8 (534)	12.0 (276)	16.4 (258)	
Alcohol intake, % (N)				
Never took alcohol	21.5 (829)	29.3 (671)	10.1 (158)	
Formerly drinking	10.8 (418)	7.78 (178)	15.3 (240)	< 0.0001
Currently drinking	67.7 (2608)	62.9 (1440)	74.6 (1168)	
Stroke, % (N)	28.9 (1123)	23.5 (541)	36.9 (582)	< 0.0001
ICV [cm ³]	1502.5 (147.4)	1422.7 (104.6)	1619.2 (120.8)	< 0.0001
TBV [cm ³]	1045.5 (98.1)	1004.7 (80.3)	1105.3 (90.8)	< 0.0001
WM [cm ³]	359.7 (44.6)	341.5 (36.6)	386.4 (41.8)	< 0.0001
Frontal GM [cm ³]	214.5 (22.1)	207.1 (19.2)	225.3 (21.7)	< 0.0001
Temporal GM [cm ³]	128.9 (13.1)	124.1 (11.1)	135.8 (12.7)	< 0.0001
Parieto-occipital GM [cm ³]	173.7 (19.0)	168.7 (16.9)	181.1 (19.4)	< 0.0001
Thalamus [cm ³]	15.1 (1.4)	14.7 (1.2)	15.8 (1.3)	< 0.0001
MTL [cm ³]	10.6 (1.1)	10.2 (1.0)	11.1 (1.1)	< 0.0001
Striatum [cm ³]	20.3 (2.3)	19.5 (2.0)	21.3 (2.2)	< 0.0001
Cerebellum [cm ³]	121.3 (12.0)	117.4 (10.6)	126.9 (11.7)	< 0.0001

BMI, body mass index; ApoE, apoE genotype; ICV, intra-cranial volume; TBV, total brain volume; WM, sum of neo-cortical white matter; Frontal GM, frontal neo-cortical gray matter; Temporal GM, temporal neo-cortical gray matter; Parieto-occipital GM, parietal and occipital neo-cortical gray matter; MTL, medial temporal lobe.

or else of otherwise stated

*

 † difference between men and women from t-test for continuous variables and chi-square test for categorical variables.

Table 2	ined mode
	-

e)	
Ģ	
0	
Ξ	
g	
e	
<u> </u>	
.Е	
q	
S	
÷	
<u> </u>	
ஆ	
بيه	
e	
U	
8	
0	
Ð	
E.	
.a	
Ŧ	
5	
-	
\mathcal{O}	
Γ	
0	
Ę	
Η	
\circ	
\sim	
-	
-	
Ę.	
nt	
ent	
erent ¹	
ferent ¹	
ifferent ¹	
different ^v	
different ¹	
of different ¹	
of different	
is of different ^v	
nts of different ^v	
ents of different ^v	
ients of different ^v	
cients of different ¹	
ficients of different ¹	
officients of different	
oefficients of different	
coefficients of different	
-coefficients of different	
ι -coefficients of different τ	
a-coefficients of different	
of α -coefficients of different ¹	
of a-coefficients of different ¹	
n of α -coefficients of different ¹	
on of α -coefficients of different '	
son of α -coefficients of different '	
rison of α -coefficients of different ¹	
arison of α -coefficients of different ¹	
parison of a-coefficients of different	
nparison of α -coefficients of different ¹	
imparison of α -coefficients of different	
comparison of α -coefficients of different	
Comparison of α-coefficients of different ¹	

				<i>p</i> -values from	comparison of a -c	coefficients		
IOA	a-coefficient (95%CI)	ΜM	Frontal GM	Temporal GM	Par-occip GM	MTL	Thalamus	Cerebellum
					All N = 3883*			
MM	1.09 (1.06–1.11)							
Frontal GM	0.740 (0.714–0.766)	<.0001						
Temporal GM	0.750 (0.724–0.774)	<.0001	0.34					
Par-occip GM	0.712 (0.683–0.741)	<.0001	0.00	0.002				
MTL	0.672 (0.642–0.702)	<.0001	<.0001	<.0001	0.02			
Thalamus	$0.588\ (0.563{-}0.613)$	<.0001	<.0001	<.0001	<.0001	<.0001		
Cerebellum	0.598 (0.570–0.627)	<.0001	<.0001	<:0001	<.0001	<.0001	0.48	
Striatum	0.552 (0.518–0.586)	<.0001	<.0001	<.0001	<.0001	<.0001	0.02	0.01
					Nomen N=2307			
WМ	1.14(1.11-1.18)							
Frontal GM	0.748 (0.709–0.787)	<.0001						
Temporal GM	0.724 (0.687–0.762)	<.0001	0.15					
Par-occip GM	0.776 (0.732–0.819)	<.0001	0.12	0.01				
MTL	$0.591\ (0.543 - 0.639)$	<.0001	<.0001	<.0001	<.0001			
Thalamus	0.590 (0.553–0.627)	<.0001	<.0001	<.0001	<.0001	0.97		
Cerebellum	0.552 (0.508–0.596)	<.0001	<.0001	<.0001	<.0001	0.18	0.11	
Striatum	0.380 (0.325–0.434)	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001
					Men N=1576			
MM	1.11 (1.07–1.16)							
Frontal GM	0.761 (0.713–0.809)	<.0001						
Temporal GM	0.742 (0.696–0.787)	<.0001	0.33					
Par-occip GM	$0.787\ (0.731 - 0.843)$	<.0001	0.24	0.07				
MTL	0.589 (0.532–0.645)	<.0001	<.0001	<.0001	<.0001			
Thalamus	0.582 (0.536-0.627)	<.0001	0.0004	<.0001	<.0001	0.81		
Cerebellum	0.523 (0.469 - 0.577)	<.0001	<.0001	<.0001	<.0001	0.07	0.04	

Author Manuscript

	Cerebellum	0.01
	Thalamus	<.0001
oefficients	MTL	<.0001
omparison of a-c	Par-occip GM	<.0001
<i>p</i> -values from c	Temporal GM	<.0001
	Frontal GM	<.0001
	ММ	<.0001
time to a see	a-coefficient (95%CI)	0.431 (0.367 - 0.495)
	101	Striatum

WM, sum of neo-cortical white matter; Frontal GM, frontal neo-cortical gray matter; Temporal GM, temporal neo-cortical gray matter; Par-occip GM, parietal and occipital neo-cortical gray matter; MTL, medial temporal lobe.

* *p*-value from log-log mixed model adjusted for age and sex. Bold figures represent non-significant p-values (> 0.00059) after Bonferroni correction.

Comparison coefficients of VOI to ICV of different quartiles of age

ION	9	-coefficients for du	ar and of age (22.20	(1)	*
101	QI [66–71 yrs]	QII [72–75 yrs]	QIII [76–79 yrs]	QIV [80–95 yrs]	d
WM	1.05 (1.01–1.09)	1.04(1.00-1.08)	1.09(1.04 - 1.14)	1.06 (1.02–1.11)	0.40
Frontal GM	0.74 (0.70–0.79)	0.71 (0.66–0.75)	0.72 (0.67–0.77)	0.66 (0.62–0.71)	0.11
Temporal GM	0.77 (0.73–0.81)	0.72 (0.68–0.76)	$0.73\ (0.68-0.78)$	0.65 (0.61–0.70)	0.004
Par-occip GM	0.71 (0.66–0.76)	0.66 (0.61–0.71)	0.69 (0.63–0.75)	0.66 (0.61–0.72)	0.50
MTL	0.67 (0.62–0.72)	0.66 (0.61–0.71)	0.63 (0.57–0.69)	0.60 (0.54–0.66)	0.32
Thalamus	0.58 (0.53–0.62)	$0.55\ (0.51{-}0.59)$	$0.56\ (0.51{-}0.61)$	$0.55\ (0.51{-}0.60)$	0.76
Striatum	0.56 (0.50-0.62)	0.52 (0.46–0.57)	$0.53\ (0.46-0.60)$	0.48 (0.42–0.55)	0.38
Cerebellum	0.58 (0.53-0.63)	0.57 (0.52–0.62)	0.57 (0.51–0.62)	0.55 (0.50-0.61)	06.0

cal gray matter; Temporal GM, temporal neo-cortical gray matter; Par- occip GM, parietal and vol, volutie of interest, yts, years, www.sunt of neo-conteat v occipital neo-cortical gray matter; MTL, medial temporal lobe.

 $_{\star}^{*}$ Linear regression with VOI as dependent variable and ICV, quartile of age and their interaction p value from the interaction term between ICV and quartile of age.

Table 4

Comparison of allometric coefficients in study populations of AGES-Reykjavik, ADNI, and CARDIA

	AGES (N=)	3883)	ADNI (N=	:180)	CARDIA (N	V=709)
101	Volume M (SD)	α. (R ²)	Volume M (SD)	α (R ²)	Volume M (SD)	α. (R ²)
ICV	1502.5 (147)	NA	1532 (156)	NA	1208 (133) *	NA
TBV	1045.5 (98)	NA	1000 (101)	NA	983 (107) *	NA
WM	359.7 (44)	1.13 (0.72)	443 (57)	1.00 (0.67)	462 (59)	1.05 (0.84)
Frontal GM	214.5 (22)	0.75 (0.52)	159 (16)	0.72 (0.59)	197 (22)	0.90 (0.76)
Temporal GM	128.9 (13)	0.73 (0.56)	122 (13)	0.81 (0.67)	156 (17)	0.91 (0.73)
Parieto-occipital GM	173.7 (19)	0.78 (0.46)	114 (12)	0.81 (0.60)	124 (14)	0.94 (0.79)
Thalamus	15.1 (1.4)	0.59 (0.39)	12 (1.4)	0.76 (0.53)	14 (1.5)	0.50 (0.26)
TTM	10.6 (1.1)	0.59 (0.43)	9.9 (1.2)	0.49 (0.31)	10 (1.1)	0.61 (0.36)
Striatum	20.3 (2.3)	0.40 (0.24)	20 (2.2)	0.62 (0.30)	20 (2.1)	$0.59\ (0.48)$
Cerebellum	121.3 (12)	$0.54\ (0.35)$	123 (12)	0.50 (0.36)	NA	NA

cranial volume; TBV = total brain volume; WM = sum of neo-cortical white matter; Frontal GM = frontal neo-cortical gray matter; Parieto-occipital GM = cranial volume; TBV = total brain volume; WM = sum of neo-cortical gray matter; Parieto-occipital GM = cranial volume; TBV = total brain volume; WM = sum of neo-cortical gray matter; Parieto-occipital GM = cranial volume; TBV = total brain volume; WM = sum of neo-cortical gray matter; Parieto-occipital GM = cranial volume; TBV = total brain volume; TBV = total brain volume; WM = sum of neo-cortical gray matter; Parieto-occipital GM = cranial volume; TBV = total brain volume; WM = sum of neo-cortical gray matter; Parieto-occipital GM = cranial volume; TBV = total brain volume; WM = sum of neo-cortical gray matter; Parieto-occipital GM = cranial volume; TBV = total brain volume; TBV = total in Young Adults Study; Volume M (SD) = mean volume in cm³ (standard deviation); α = allometric coefficient derived from [log (VOI) = intercept + α log (VOI) + β age × age + β sex × sex]; ICV = intratroimaging Initiative; CARDIA = Coronary Artery Risk Development parietal and occipital neo-cortical gray matter; MTL = medial temporal lobe. à

* Includes supratentorial areas only