

- 1) Wolk, D. A., Das, S. R., Mueller, S. G., Weiner, M. W., & Yushkevich, P. A. (2017). Medial temporal lobe subregional morphometry using high resolution MRI in Alzheimer's disease. *Neurobiology of Aging*, 49. <https://doi.org/10.1016/j.neurobiolaging.2016.09.011>
- 2) Mueller, S. G., Yushkevich, P. A., Das, S., Wang, L., Van Leemput, K., Iglesias, J. E., ... Weiner, M. W. (2018). Systematic comparison of different techniques to measure hippocampal subfield volumes in ADNI2. *NeuroImage: Clinical*, 17. <https://doi.org/10.1016/j.nicl.2017.12.036>
- 3) Das, S.R., Xie, L., Wisse, L.E.M., Vergnet, N., Ittyerah, R., Cui, S., Yushkevich, P.A., and Wolk, D.A. for the Alzheimer's Disease Neuroimaging Initiative (2019). In-vivo Measures of Tau Burden are Associated with Atrophy in Early Braak Stage Medial Temporal Lobe Regions in Amyloid Negative Individuals. *Alzheimer's and Dementia*, in press.

Outside lab using ADNI T2 data:

- 1) Iglesias, J. E., Van Leemput, K., Augustinack, J., Insausti, R., Fischl, B., Reuter, M., & Alzheimer's Disease Neuroimaging Initiative. (2016). Bayesian longitudinal segmentation of hippocampal substructures in brain MRI using subject-specific atlases. *NeuroImage*, 141, 542–555. <https://doi.org/10.1016/j.neuroimage.2016.07.020>